



# **Mark Scheme (Results)**

Summer 2017

Pearson Edexcel International GCSE  
in Chemistry (4CH0) Paper 1C  
Science (Double Award) (4SC0) Paper 1C

Pearson Edexcel Level 1/Level 2 Certificate  
Biology (KCH0) Paper 1C  
Science (Double Award) (KSC0) Paper 1C

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2017

Publications Code 4CH0\_1C\_1706\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2017

## **General Marking Guidance**

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

| Question number                                    | Answer  | Notes                     | Marks |                        |   |                         |   |   |   |  |   |  |   |
|--|---|---------------------------|-------|------------------------|---|-------------------------|---|---|---|--|---|--|---|
| 1  | <table border="1"> <tr> <td data-bbox="416 186 931 244">atomic number of the atom</td> <td data-bbox="931 186 1128 244">3</td> </tr> <tr> <td data-bbox="416 244 931 302">number of shells shown</td> <td data-bbox="931 244 1128 302">2</td> </tr> <tr> <td data-bbox="416 302 931 361">mass number of the atom</td> <td data-bbox="931 302 1128 361">7</td> </tr> <tr> <td data-bbox="416 361 931 455">number of protons in an isotope of this element</td> <td data-bbox="931 361 1128 455">3</td> </tr> <tr> <td data-bbox="416 455 931 550">group where element is found in the Periodic Table</td> <td data-bbox="931 455 1128 550">1</td> </tr> </table> | atomic number of the atom | 3     | number of shells shown | 2 | mass number of the atom | 7 | number of protons in an isotope of this element | 3 | group where element is found in the Periodic Table | 1 |  | 5 |
| atomic number of the atom                          | 3   |                           |       |                        |   |                         |   |   |   |  |   |  |   |
| number of shells shown                             | 2   |                           |       |                        |   |                         |   |   |   |  |   |  |   |
| mass number of the atom                            | 7   |                           |       |                        |   |                         |   |   |   |  |   |  |   |
| number of protons in an isotope of this element    | 3   |                           |       |                        |   |                         |   |   |   |  |   |  |   |
| group where element is found in the Periodic Table | 1   |                           |       |                        |   |                         |   |   |   |  |   |  |   |

**(Total for Question 1 = 5 marks)**

| Question number | Answer   | Notes | Marks |
|-----------------|--|-------|-------|
| 2 (a)           | <p><b>B (H<sub>2</sub>)</b></p> <p><b>The only correct answer is B</b></p> <p>A is not correct because H is the symbol of an atom of hydrogen not a molecule of hydrogen</p> <p>C is not correct because H<sub>2</sub>O is the formula of water which is a compound</p> <p>D is not correct because H<sub>2</sub>O<sub>2</sub> is the formula of hydrogen peroxide which is a compound</p> |       | 1     |
| (b)             | <p><b>D (sodium chloride solution)</b></p> <p><b>The only correct answer is D</b></p> <p>A is not correct because sodium is an element not a mixture</p> <p>B is not correct because chlorine is an element not a mixture</p> <p>C is not correct because sodium chloride is a pure compound not a mixture</p>   |       | 1     |

| Question number | Answer   | Notes  | Marks |
|-----------------|--|--|-------|
| (c)             | <p><b>A (chromatography)</b></p> <p><b>The only correct answer is A</b></p> <p>B is not correct because crystallisation would not separate the dyes in food colouring</p> <p>C is not correct because evaporation would not separate the dyes in food colouring</p> <p>D is not correct because filtration would not separate the dyes in food colouring</p> |  | 1     |
| (d) (i)         | <p>M1 add (dilute) nitric acid</p><br><p>M2 add silver nitrate (solution)</p><br><p>M3 white precipitate</p>   | <p>Accept HNO<sub>3</sub></p> <p>If no acid then M2 M3 can be scored</p> <p>If incorrect acid then M2 M3 can be scored</p><br><p>Ignore references to testing for chlorine/electrolysis</p><br><p>Accept AgNO<sub>3</sub></p> <p>Do not award mark if additional reagent added</p> <p>Ignore litmus</p><br><p>Accept usual alternatives to precipitate</p> <p>Ignore cloudy/milky</p> <p>Ignore qualifiers such as pale</p> <p>Reject all other colours</p> <p>Reject other observations e.g. fizzing</p><br><p>M3 DEP on mention of silver nitrate/AgNO<sub>3</sub></p> | 3     |

| Question number | Answer  | Notes                          | Marks |
|-----------------|---|--------------------------------|-------|
| (d) (ii)        | diffusion   | Ignore identity of precipitate | 1     |
| (e) (i)         | M1 flask<br>M2 column<br>M3 condenser   |                                | 3     |
| (ii)            | <p><b>C (g) → (l)</b></p> <p><b>The only correct answer is C</b></p> <p>A is not correct because change of state from solid to aqueous does not occur in fractional distillation</p> <p>B is not correct because change of state from liquid to solid does not occur in fractional distillation</p> <p>D is not correct because change of state from aqueous to solid does not occur in fractional distillation</p> |                                | 1     |

**(Total for Question 2 = 11 marks)**

| Question number | Answer                    | Notes  | Marks |
|-----------------|---------------------------|--|-------|
| 3 (a)           | galvanising/galvanisation | Ignore sacrificial protection  | 1     |
| (b) (i)         | rust                      | Accept air / O <sub>2</sub><br>Ignore O<br><br>Accept H <sub>2</sub> O<br>Allow moisture<br>Ignore vapour<br><br>Accept answers in either order<br>If name and formulae given both must be correct | 1     |
|                 | M1 oxygen<br><br>M2 water |  | 2     |



| Question number | Answer  | Notes  | Marks |
|-----------------|---|--|-------|
| (iii)           | <p>M1 Zn/zinc is more reactive than Fe/iron</p> <p>M2 Zn loses (two) electrons / Zn forms Zn<sup>2+</sup></p> <p>M3 prevents Fe from losing electrons / prevents Fe forming Fe<sup>2+</sup></p> | <p>Accept Zn higher in reactivity series<br/> Accept zinc reacts in preference to/instead of iron<br/> Allow zinc is oxidised in preference to iron<br/> Ignore references to sacrificial protection<br/> Reject zinc rusts<br/> Reject use of Zn<sup>2+</sup> and Fe<sup>2+</sup> in place of Zn / Fe</p> <p>Accept reverse argument for iron</p> <p>Accept Zn → Zn<sup>2+</sup> (+ 2e<sup>-</sup>)<br/> Reject Fe loses (two) electrons/Fe forms Fe<sup>2+</sup></p> <p>Accept (so) Fe → Fe<sup>2+</sup> (+ 2e<sup>-</sup>) does not take place</p> <p>Accept (so) Fe<sup>2+</sup> (+ 2e<sup>-</sup>) → Fe (does take place) IF mentioned in correct context of reaction with zinc/displacement</p> <p>Accept Fe<sup>2+</sup> (ions) gain (two) electrons and converted into Fe (atoms) IF mentioned in context of reaction with zinc/displacement</p> | 3     |

**(Total for Question 3 = 7 marks)**

| Question number | Answer  | Notes | Marks |
|-----------------|---|-------|-------|
| 4 (a)           | M1 (top row) $\text{CuCl}_2$<br>M2 (middle row) $(\text{NH}_4)_2\text{SO}_4$<br>M3 (bottom row) $\text{Fe}_2(\text{CO}_3)_3$  |       | 3     |
| (b)             | <p><b>C (copper(II) sulfate)</b></p> <p><b>The only correct answer is C</b></p> <p>A is not correct because the name of <math>\text{CuSO}_4</math> is not copper(I) sulfate</p> <p>B is not correct because the name of <math>\text{CuSO}_4</math> is not copper(I) sulfite</p> <p>D is not correct because the name of <math>\text{CuSO}_4</math> is not copper(II) sulphite</p> |       | 1     |
| (c)             | <p><b>C (white AND colourless)</b></p> <p><b>The only correct answer is C</b></p> <p>A is not correct because <math>\text{NH}_4\text{Cl}(s)</math> is not colourless</p> <p>B is not correct because <math>\text{NH}_4\text{Cl}(s)</math> is not colourless</p> <p>D is not correct because <math>\text{NH}_4\text{Cl}(aq)</math> is not white</p>                                |       | 1     |

| Question number | Answer   | Notes  | Marks |
|-----------------|--|--|-------|
| (d) (i)         | <p><b>A (brown precipitate)</b></p> <p><b>The only correct answer is A</b></p> <p>B is not correct because the product of the test, iron(III) hydroxide is not a brown solution</p> <p>C is not correct because the product of the test, iron(III) hydroxide is not a green precipitate</p> <p>D is not correct because the product of the test, iron(III) hydroxide is not a green solution</p> |  | 1     |
| (ii)            | <p>M1 iron(III) hydroxide</p> <p>M2 sodium sulfate</p>   | <p>Ignore iron hydroxide<br/>Accept ferric hydroxide<br/>Ignore formulae whether correct or incorrect</p> <p>Accept sulphate</p> <p>Accept answers in either order</p>                           | 2     |
| (iii)           | <p>to prevent the formation of other precipitates<br/>OR<br/>to react with/remove carbonate (ions)</p>   | <p>Ignore references to impurities</p> <p>Accept so only sulfate (ions) react with barium chloride/barium ions</p> <p>Accept to remove sulfite ions<br/>Reject if to remove an incorrect ion</p> | 1     |

| Question number | Answer  | Notes   | Marks |
|-----------------|---|---|-------|
| (d) (iv)        | white precipitate   | Reject other colours<br>Accept usual alternatives for precipitate<br>Reject incorrect extra observations  | 1     |
| (e)             | M1 add (dilute) acid /H <sup>+</sup><br><br>OR heat<br><br>M2 bubble/pass gas/carbon dioxide into limewater/OWTTE<br><br>M3 (limewater) turns milky | Accept any named acid<br>Accept correct formulae<br><br>Reject if limewater added to CuCO <sub>3</sub><br><br>Accept cloudy / white precipitate<br>M3 indep | 3     |

**(Total for Question 4 = 13 marks)**

| Question number | Answer   | Notes   | Marks |
|-----------------|--|---|-------|
| 5 (a) (i)       | to (produce) heat (energy)   | Ignore exothermic reaction<br>Accept to reach/maintain a high temperature<br>Allow to make the furnace/it hot<br>Ignore to increase the temperature<br>Ignore for energy alone<br>Ignore to make carbon dioxide | 1     |
| (ii)            | to produce the reducing agent  | Accept to produce substance needed to reduce iron ore/iron oxide/haematite<br><br>Ignore to make carbon monoxide<br>Allow to make carbon monoxide/CO/substance which reacts with iron ore to produce iron       | 1     |
| (b)             | M1 $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$<br>M2 $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$ | Award 1 mark for two correct equations in wrong order<br>Ignore state symbols   | 2     |

| Question number | Answer   | Notes   | Marks |
|-----------------|--|---|-------|
| (c)             | <p>M1 iron/iron oxide is reduced AND carbon is oxidised</p> <p>M2 (because) iron/iron oxide loses oxygen AND carbon gains oxygen</p> | <p>In M1 and M2 Accept Fe/Fe<sub>2</sub>O<sub>3</sub>/Fe<sup>3+</sup> and C</p> <p>Allow use of O</p> <p>Ignore references to electrons</p> <p>Allow reference to oxidation number of iron has decreased and oxidation number of carbon has increased (if oxidation numbers stated must be correct)</p> | 2     |

**(Total for Question 5 = 6 marks)**

| Question number  | Answer   | Notes   | Marks |  |  |   |   |   |  |   |   |  |  |   |   |
|--|--|---|-------|--|--|---|---|---|--|---|---|--|--|---|---|
| 6 (a)  | M1 vapourisation<br>M2 temperature<br>M3 condensation<br>M4 boiling point  |   | 4     |  |  |   |   |   |  |   |   |  |  |   |   |
| (b)  | <table border="1"> <tbody> <tr> <td data-bbox="434 500 994 598">the molecules that are cracked are hydrocarbons</td> <td data-bbox="994 500 1167 598">✓</td> </tr> <tr> <td data-bbox="434 598 994 695">catalytic cracking uses iron as the catalyst</td> <td data-bbox="994 598 1167 695"></td> </tr> <tr> <td data-bbox="434 695 994 793">cracking is used because of different requirements for hydrocarbons</td> <td data-bbox="994 695 1167 793">✓</td> </tr> <tr> <td data-bbox="434 793 994 928">cracking reactions are examples of addition reactions</td> <td data-bbox="994 793 1167 928"></td> </tr> <tr> <td data-bbox="434 928 994 1064">cracking produces molecules with shorter chains</td> <td data-bbox="994 928 1167 1064">✓</td> </tr> <tr> <td data-bbox="434 1064 994 1199">CH<sub>4</sub> + 2O<sub>2</sub> → CO<sub>2</sub> + 2H<sub>2</sub>O is an equation for a cracking reaction</td> <td data-bbox="994 1064 1167 1199"></td> </tr> </tbody> </table> | the molecules that are cracked are hydrocarbons | ✓     | catalytic cracking uses iron as the catalyst |  | cracking is used because of different requirements for hydrocarbons | ✓ | cracking reactions are examples of addition reactions |  | cracking produces molecules with shorter chains | ✓ | CH <sub>4</sub> + 2O <sub>2</sub> → CO <sub>2</sub> + 2H <sub>2</sub> O is an equation for a cracking reaction |  | Award 1 mark for each correct tick<br>If more than 3 ticks then subtract 1 mark for each extra tick | 3 |
| the molecules that are cracked are hydrocarbons  | ✓  |   |       |  |  |   |   |   |  |   |   |  |  |   |   |
| catalytic cracking uses iron as the catalyst   |  |   |       |  |  |   |   |   |  |   |   |  |  |   |   |
| cracking is used because of different requirements for hydrocarbons  | ✓  |   |       |  |  |   |   |   |  |   |   |  |  |   |   |
| cracking reactions are examples of addition reactions  |  |   |       |  |  |   |   |   |  |   |   |  |  |   |   |
| cracking produces molecules with shorter chains  | ✓  |   |       |  |  |   |   |   |  |   |   |  |  |   |   |
| CH <sub>4</sub> + 2O <sub>2</sub> → CO <sub>2</sub> + 2H <sub>2</sub> O is an equation for a cracking reaction |  |   |       |  |  |   |   |   |  |   |   |  |  |   |   |

| Question number | Answer  | Notes  | Marks |
|-----------------|---|--|-------|
| 6 (c) (i)       | $C_nH_{2n}$   | Accept other letters, such as x, in place of n   | 1     |
| (ii)            | M1 propene<br>M2 methane  | Accept propylene<br>Accept answers in either order   | 2     |
| (iii)           | <pre>       H H           H - C - C - H                 H H </pre>  | Must show all atoms and all bonds  | 1     |
| (iv)            | <pre>       H H H H               H - C = C - C - C - H                                 H H        H H H H               H - C - C = C - C - H                           H           H </pre> | Award 1 mark for each structure<br>Accept answers in either order<br>Must show all atoms and all bonds<br><br>Allow cis/trans isomers<br>Allow cyclobutane | 2     |



| Question number | Answer   | Notes  | Marks |
|-----------------|--|--|-------|
| (d)             | <p>M1 product has longer (carbon) chain</p> <p>M2 only product has (all) single (C—C) bonds</p> <p>M3 only the product is a solid / only the reactant is a gas</p> | <p>Must have a comparative statement / a statement about both reactant and product</p> <p>Accept reactant has shorter (carbon) chain</p> <p>Ignore reactant is a monomer and product is a polymer</p> <p>Accept only the reactant has a double bond<br/>Allow only the reactant is unsaturated<br/>Allow only the product is saturated</p> <p>Ignore references to alkanes/alkene</p> <p>Reject if an incorrect state also given</p> | 3     |


**(Total for Question 6 = 16 marks)**

| Question number | Answer  | Notes   | Marks |
|-----------------|---|---|-------|
| 7 (a)           | reaction faster / takes less time                       | Ignore references to larger surface area<br>Reject references to dissolving   | 1     |
| (b)             | reaction very slow/too slow without water               | Accept rusting needs water/moisture<br>Allow so that iron can rust<br>Accept reference to increasing rate of reaction/rusting | 1     |
| (c)             | M1 (at start) 71<br>M2 (at end) 16<br>M3 (change) (-)55 | Ignore trailing .0(0) in all answers<br>Ignore units<br><br>M3 CQ M1 -M2  | 3     |

| Question number | Answer   | Marks |
|-----------------|--|-------|
| (d)             | <p><b>D (he uses a smaller mass of iron filings)</b><br/><b>The only correct answer is D</b></p> <p>A is not correct because the change in volume is governed by the mass of iron so to produce a smaller change in volume the mass of iron present needs to be smaller – increasing the volume of water would not produce a smaller volume</p> <p>B is not correct because the change in volume is governed by the mass of iron so to produce a smaller change in volume the mass of iron present needs to be smaller – leaving the apparatus for longer would not produce a smaller change in volume</p> <p>C is not correct because the change in volume is governed by the mass of iron so to produce a smaller change in volume the mass of iron present needs to be smaller – having the apparatus in a warmer place would change the rate of reaction – and would not affect the change in volume</p> | 1     |

| Question number | Answer   |  | Marks |
|-----------------|--|--|-------|
| (e) (i)         | identifies the error e.g. has divided by the syringe reading at the start<br><br>OR<br><br>corrects the error e.g. should divide by the total volume | Accept has divided by 90 / has divided by the wrong number/has not included the volume of air in the flask and glass tube/has not included the 250<br><br>Accept should divide by 340 / should include the volume of air in the flask and glass tube | 1     |
| (ii)            | M1 $\frac{70 \times 100}{340}$ (= 20.5882.....)<br><br>M2 21   | Allow any number of sig. fig. but reject 20  | 2     |
|                 |  | Allow $\frac{70 \times 100}{250} = 28$ for (1)<br><br>Correct final answer with or without working scores 2 marks  |       |

**(Total for Question 7 = 9 marks)**

| Question number   | Answer   | Notes   | Marks      |
|-------------------|--|---|------------|
| 8 (a) (i)<br>(ii) |   | <p>curve from reactant level to product level with peak below that of original (1)</p> <p>M1 for approximately vertical line/arrow between reactant level and product level labelled <math>\Delta H</math>/enthalpy change/<math>-9 \text{ kJ/mol}</math></p> <p>M2 for approximately vertical line/arrow between reactant level and peak of <u>candidate</u> curve labelled <math>E_{\text{cat}}</math> /activation energy</p> <p>M1 and M2 CQ candidate curve</p> | 1<br><br>2 |
| (b) (i)           | rate decreases / OWTTE   | <p>Allow (reaction is) slower</p> <p>Allow reaction takes longer</p> <p>Ignore references to yield / position of equilibrium</p>  | 1          |
| (ii)              | <p>M1 (at lower temperature equilibrium position shifts to right so yield of hydrogen iodide) increases</p> <p>M2 because (forward) reaction is exothermic</p> | <p>Ignore <math>\Delta H</math> is negative</p> <p>Accept backward reaction is endothermic</p> <p>Ignore because reaction moves in exothermic direction</p> <p>Ignore references to Le Chatelier's principle e.g. decrease in temperature favours exothermic reaction</p> <p>M2 DEP M1</p>  | 2          |

| Question number | Answer   | Notes   | Marks |
|-----------------|--|---|-------|
| (c) (i)         | (rate) decreases / OWTTE   | Allow (reaction is) slower<br>Allow reaction takes longer (to reach equilibrium)<br><br>Ignore references to yield / position of equilibrium  | 1     |
| (ii)            | M1 (decrease in pressure has) no effect (on yield of hydrogen iodide)<br><br>M2 because equal numbers of (gas) moles/molecules on both sides | Allow no change<br>Ignore has no effect on other factors e.g. equilibrium (position)<br>Ignore references to rate<br><br>Allow (gas) particles for moles/molecules<br><br>M2 DEP M1 | 2     |

**(Total for Question 8 = 9 marks)**

| Question number  | Answer  | Notes  | Marks |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
|--|---|--|-------|--|--|--|--|--|---|-----------------------------------|---|-----------------------------------|--|---|---|
| 9 (a)  | <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px;">the elements can be obtained by electrolysing molten metal halides</td> <td style="text-align: center; width: 50px;">✓</td> </tr> <tr> <td style="padding: 5px;">the elements with paler colours are lower down the group</td> <td></td> </tr> <tr> <td style="padding: 5px;">the boiling points decrease down the group</td> <td></td> </tr> <tr> <td style="padding: 5px;">the elements form covalent compounds with other non-metals</td> <td style="text-align: center;">✓</td> </tr> <tr> <td style="padding: 5px;">their molecules contain two atoms</td> <td style="text-align: center;">✓</td> </tr> <tr> <td style="padding: 5px;">all are gases at room temperature</td> <td></td> </tr> </table> | the elements can be obtained by electrolysing molten metal halides | ✓     | the elements with paler colours are lower down the group |  | the boiling points decrease down the group |  | the elements form covalent compounds with other non-metals | ✓ | their molecules contain two atoms | ✓ | all are gases at room temperature |  | Award 1 mark for each correct tick<br>If more than 3 ticks then subtract 1 mark for each extra tick | 3 |
| the elements can be obtained by electrolysing molten metal halides | ✓   |  |       |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
| the elements with paler colours are lower down the group           |   |  |       |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
| the boiling points decrease down the group                         |   |  |       |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
| the elements form covalent compounds with other non-metals         | ✓   |  |       |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
| their molecules contain two atoms                                  | ✓   |  |       |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
| all are gases at room temperature                                  |   |  |       |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
| (b)  | $2K + I_2 \rightarrow 2KI$  | Accept fractions/multiples<br>Ignore state symbols                 | 1     |  |  |  |  |  |   |                                   |   |                                   |  |   |   |
| (c)  | (hydrogen chloride)      g<br>(hydrochloric acid)      aq   |  | 2     |  |  |  |  |  |   |                                   |   |                                   |  |   |   |

| Question number   | Answer   | Notes   | Marks |
|---|--|---|-------|
| (d)   | M1 effervescence / bubbles / fizzing   | Accept gas formed / evolved / given off                             | 3     |
|   |  | Ignore hydrogen   |       |
|   |  | Reject extra incorrect observations                                 |       |
|   | M2 (in water hydrogen chloride forms) H <sup>+</sup> ions / forms (hydrochloric) acid  | Ignore dissociates/ionises if no reference to H <sup>+</sup> / acid |       |
|   | M3 magnesium reacts to form hydrogen/H <sub>2</sub>  | Accept chemical/word equation<br>M3 DEP M2                          |       |
|   | <b>OR</b>  |   |       |
| M1 magnesium disappears/gets smaller  | Allow dissolves<br>Reject extra incorrect observations<br>Ignore magnesium moving  |   |       |
| M2 (in water hydrogen chloride forms) H <sup>+</sup> ions / forms (hydrochloric) acid             | Ignore dissociates/ionises if no reference to H <sup>+</sup> / acid  |   |       |
| M3 magnesium reacts to form magnesium chloride/MgCl <sub>2</sub> /magnesium ions/Mg <sup>2+</sup> | Accept chemical/word equation<br>M3 DEP M2<br><br>Ignore references to solution with HCl dissolved in methylbenzene before water added |   |       |



| Question number | Answer  | Notes  | Marks |
|-----------------|---|--|-------|
| 9 (e) (i)       | M1 test 2<br>M2 chlorine does not react with chloride (ions)  | Allow description of test<br>Accept chlorine does not displace itself<br>Allow chlorine does not react with itself<br>Accept reference to halogen/halide in place of chlorine/chloride<br>Ignore chlorine does not displace chloride<br>M2 DEP on correct M1 or missing M1 | 2     |
| (ii)            | <p><b>C (solution becomes darker)</b></p> <p><b>The only correct answer is C</b></p> <p>A is not correct because original sodium iodide solution is colourless but in Test 1 iodine is formed in the solution so the solution becomes darker (red/brown)- no effervescence is seen as no gas being formed</p> <p>B is not correct because original sodium iodide solution is colourless but in Test 1 iodine is formed in the solution so the solution becomes darker (red/brown)- no iodine vapour produced so no purple fumes produced</p> <p>D is not correct because original sodium iodide solution is colourless but in Test 1 iodine is formed in the solution so the solution becomes darker (red/brown)- no white precipitate formed</p> |  | 1     |

| Question number | Answer  | Notes  | Marks |
|-----------------|---|--|-------|
| (f) (i)         | $\text{Cl}_2 + 2\text{At}^- \rightarrow \text{At}_2 + 2\text{Cl}^-$   | Accept fractions/multiples<br>Ignore 2e on both sides of equation<br>Reject other extra species  | 1     |
| (ii)            | M1 chlorine/ $\text{Cl}_2$ is reduced AND astatide (ion)/ $\text{At}^-$ is oxidised<br><br>M2 chlorine/ $\text{Cl}_2$ gains electron(s) AND astatide (ion)/ $\text{At}^-$ loses electron(s) | Allow Cl<br>Reject chloride (ions) reduced<br><br>Allow Cl<br>Ignore references to oxidation numbers<br><br>Reject use of astatine in place of astatide ions once only in (ii) | 2     |

**(Total for Question 9 = 15 marks)**

| Question number | Answer   | Notes  | Marks |
|-----------------|--|--|-------|
| 10 (a)          | <p>M1 replace — signs by + signs / OWTTE</p> <p>M2 replace protons (label) by electrons</p>  | <p>Accept all ions should be cations/positive<br/>Reject if state/imply these particles are protons/nuclei</p> <p>Allow electrons (not protons) are delocalised</p>  | 2     |
| (b)             | <p>M1 (magnesium) ions in layers/rows/sheets/planes/OWTTE</p> <p>M2 slip / slide (over each other)</p> <p>M3 delocalised electrons</p> <p>M4 can flow/move (through the magnesium/metal/structure) / are mobile (when voltage/p.d. is applied)</p> | <p>Accept atoms/cations/particles for ions<br/>Reject molecules</p> <p>Allow OWTTE eg shift/roll/flow</p> <p>M2 DEP on mention of EITHER layers or equivalent<br/>OR mention of ions or equivalent</p> <p>Do not award M2 if molecules/protons/electrons/nuclei in place of ions etc.</p> <p>If reference to ionic bonding / covalent bonding / molecules / intermolecular forces, M1 and M2 cannot be scored</p> <p>Accept sea of electrons<br/>Ignore free electrons</p> <p>Ignore carry the charge<br/>M4 DEP on mention of electrons</p> | 4     |

| Question number | Answer   | Notes  | Marks |
|-----------------|--|--|-------|
| (c)<br>(i)      | M1    bright / white flame<br><br>M2    white solid / powder / ash / smoke   | Allow bright / white light<br>Ignore sparks<br><br>Ignore grey<br>Reject precipitate | 2     |
| (ii)            | <p><b>A    (it makes a squeaky pop with a lighted splint)</b></p> <p><b>The only correct answer is A</b></p> <p>B is not correct because hydrogen gas does not relight a glowing splint</p> <p>C is not correct because hydrogen gas does not turn damp blue litmus red</p> <p>D is not correct because hydrogen gas does not turn limewater milky</p> |  | 1     |

| Question number | Answer   | Notes  | Marks |
|-----------------|--|--|-------|
| (d)             | to evaporate/remove/boil off some of the water   | Reject to remove (all) the water (stated or implied)<br><br>Accept to increase the concentration (of the solution)<br>Accept to produce a (hot) saturated solution<br>Accept to reach crystallisation point<br>Allow to reduce the volume (of the solution)<br>Allow to evaporate/remove some of the solution<br>Ignore references to rate of reaction<br>Reject to evaporate acid | 1     |
|                 | M1 to see if/when crystals form  | Allow solid for crystals   | 2     |
|                 | M2 indicates the crystallisation point/when solution is saturated/OWTTE                          | Accept indicates when to stop heating/if need to continue heating  |       |
|                 | Any two from<br>M1 MgSO <sub>4</sub><br>M2 H <sub>2</sub> O<br>M3 H <sub>2</sub> SO <sub>4</sub> | Ignore names even if incorrect   | 2     |

| Question number | Answer  | Notes  | Marks |
|-----------------|---|--|-------|
| 10 (e)          | <p>M1 calculation of mass of water</p> <p>M2 calculation of amounts of MgSO<sub>4</sub> AND H<sub>2</sub>O</p> <p>M3 calculation of MgSO<sub>4</sub> : H<sub>2</sub>O ratio</p> <p>M4 x=7</p> | <p><math>m(\text{H}_2\text{O}) = (17.2 - 8.3) = 8.9 \text{ (g)}</math></p> <p><math>n(\text{MgSO}_4) = 8.3 \div 120 \text{ OR } 0.069 \text{ (mol)}</math><br/>Allow any sig. fig e.g. 0.07</p> <p>AND</p> <p><math>n(\text{H}_2\text{O}) = 8.9 \div 18 \text{ OR } 0.49 \text{ (mol)}</math><br/>Allow any sig. fig e.g. 0.5</p> <p>0.069 : 0.49 OR 1 : 7.101449275..<br/>allow any sig. fig.</p> <p>OR 0.49/0.069 OR 7.101449275.. allow any sig. fig.</p> <p>accept MgSO<sub>4</sub>.7H<sub>2</sub>O</p> <p>Correct final answer with or without working scores 4 marks</p> <p>Check for incorrect working<br/>e.g. <math>120 \div 18 = 6.66 = 7</math></p> <p>Mark CQ throughout</p> | 4     |

**(Total for Question 10 = 18 marks)**

| Question number | Answer  | Notes   | Marks |
|-----------------|---|---|-------|
| 11 (a) (i)      | <p>(moles route)</p> <p>M1 <math>n(\text{CaO}) = 28000 \div 56</math> OR 500 (mol)</p> <p>M2 <math>n(\text{H}_2\text{O}) = 500</math> (mol)</p> <p>M3 <math>m(\text{H}_2\text{O}) = 500 \times 18 = 9000</math> g / 9 kg</p> <p>OR<br/>(mass ratios route)</p> <p>M1 <math>M_r(\text{CaO}) = 56</math> AND <math>M_r(\text{H}_2\text{O}) = 18</math></p> <p>M2 <math>m(\text{H}_2\text{O}) = \frac{28 \times 18}{56}</math></p> <p>M3 = 9 kg / 9000 g</p> | <p>Allow 0.5 if final answer given in kg</p> <p>For M2 need to use or state mol H<sub>2</sub>O</p> <p>Units not needed for intermediate answers</p> <p>Do not award M3 if unit missing or incorrect</p> <p>Correct final answer with or without working scores 3 marks</p> <p>Mark M2 and M3 CQ on M1</p> <p>Do not award M3 if unit missing or incorrect</p> <p>Correct final answer with or without working scores 3 marks</p> <p>Mark M2 and M3 CQ on M1</p> | 3     |
| (ii)            | <p>M1 carbon dioxide is (an) acidic (oxide)</p> <p>M2 calcium hydroxide is a base / an alkali</p>   | <p>Ignore contains hydroxide/OH<sup>-</sup> ions</p>  | 2     |

| Question number | Notes   | Marks |
|-----------------|---|-------|
| 11 (b)          | <p>M1 + M2 all six points plotted correctly<br/>Deduct 1 mark for each error up to max 2</p> <p>M3 curve of best fit<br/>Curve CQ on points plotted<br/>Penalise repeated straight line(s) joining points<br/>Penalise more than one curve visible</p>  | 3     |
| (c)             | <p>Ignore move faster</p> <p>Allow more molecules/particles have enough energy for successful collisions OWTTE</p> <p>Ignore references to probability /chance/likelihood of collisions</p> <p>M3 DEP on mention of energy in M1 or M2</p> <p>Penalise use of atoms/ions instead of molecules/particles once only</p> | 3     |

**(Total for Question 11 = 11 marks)**



Pearson Education Limited. Registered company number 872828  
with its registered office at 80 Strand, London WC2R 0RL