



Mark Scheme (Results)

Summer 2018

Pearson Edexcel GCE
In Chemistry (8CH0) Paper 02
Core Organic and Physical Chemistry

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2018

Publications Code 8CH0_02_1806_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question Number	Acceptable Answer	Mark
1	<p data-bbox="362 268 837 304">The only correct answer is A</p> <p data-bbox="362 339 1234 376"><i>B is incorrect because H₂O is a nucleophile – via lone pairs</i></p> <p data-bbox="362 411 1227 448"><i>C is incorrect because NH₃ is a nucleophile – via lone pair</i></p> <p data-bbox="362 483 1234 520"><i>D is incorrect because CN⁻ is a nucleophile – via lone pairs</i></p>	(1)

(Total for Question 1 = 1 mark)

Question Number	Acceptable Answer	Additional Guidance	Mark
2 (a)(i)	<ul style="list-style-type: none"> converts temperature to Kelvin and pressure to Nm^{-2} (Pa) (1) rearranging ideal gas equation and substituting their values (1) evaluates answer to 2 SF and includes units (1) 	<p><u>Examples of calculation</u> $60^\circ\text{C} = 333 \text{ K}$ $500 \text{ kPa} = 5 \times 10^5 / 500\,000 \text{ Pa}$</p> $V = \frac{nRT}{P}$ $V = 1 \times 8.31 \times 333 / 500\,000$ $= 5.53446 \times 10^{-3}$ $= 0.0055 \text{ m}^3 / 5.5 \times 10^{-3} \text{ m}^3 / 5.5 \text{ dm}^3 / 5500 \text{ cm}^3$ <p>allow TE answers to 2 SF only correct answer with no working scores 3 marks correct answer with incorrect working scores 2 marks max.</p>	(3)

Question Number	Acceptable Answer	Additional Guidance	Mark
2(a)(ii)	<ul style="list-style-type: none"> calculates M_r to 2 or more SF (1) identifies element X (1) 	<p><u>Example of calculation:</u> molar mass = mass in 24000 cm^3 $= 1.42 \times 24000 / 1000 = 34 (.08) \text{ (g mol}^{-1}\text{)}$ ignore SF except 1 SF</p> $(X + (3 \times 1)) = 34$ $X = 31 \text{ so P / phosphorus}$ <p>just 'phosphorus' with no working scores M2 only</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
2(b)(i)	<ul style="list-style-type: none"> • calculates moles of acid (1) • calculates moles of sodium carbonate (1) • recognises that (sodium) carbonate is in excess (1) • evidence for excess sodium carbonate in terms of moles (1) • correct volume of gas calculated with units (1) 	<p><u>Example of calculation</u></p> <p>moles of acid = $10.0 \times 0.400/1000$ = $4(.0) \times 10^{-3}/0.004$ (mol)</p> <p>moles of sodium carbonate = $0.242/106.0$ = $2.283 \times 10^{-3}/0.002283$ (mol)</p> <p>recognition of $\text{HCl}:\text{Na}_2\text{CO}_3 = 2:1$ gets M4 4.0×10^{-3} mol acid requires 2.0×10^{-3} mol sodium carbonate OR 2.283×10^{-3} mol of sodium carbonate requires 4.566×10^{-3} mol of acid</p> <p>moles $\text{CO}_2 = 2.0 \times 10^{-3}$ (mol)</p> <p>volume of gas = $2.0 \times 10^{-3} \times 24\,000$ = $48 \text{ cm}^3/0.048 \text{ dm}^3$ TE on incorrect moles CO_2 correct answer with no working scores 1 mark if the moles of sodium carbonate are not calculated, only M1, M4 and M5 can be awarded. ignore SF except 1 for M5</p>	(5)

Question Number	Acceptable Answer	Additional Guidance	Mark
2(b)(ii)	<p>An answer that makes reference to the following reasons:</p> <ul style="list-style-type: none"> • some gas escaped before the bung/delivery tube was replaced (1) • the gas / carbon dioxide is (slightly) soluble in water/ acid / solution (1) 	<p>ignore references to change in volume when the bung is pushed into the test tube</p> <p>allow 'temperature less than 25°C/298 K/room temperature' as alternative to either answer</p> <p>do not award an incomplete reaction</p> <p>do not award leaky apparatus/sticking syringe</p>	(2)

(Total for Question 2 = 12 marks)

Question Number	Acceptable Answer	Additional Guidance	Mark
3(a)	<ul style="list-style-type: none"> K_c expression 	(1) $(K_c =) \frac{[N_2(g)]^2 [H_2O(g)]^6}{[NH_3(g)]^4 [O_2(g)]^3}$	(2)
	<ul style="list-style-type: none"> units based on their K_c expression 	(1) ignore missing state symbols do not award round brackets mol dm ⁻³ or mol/dm ³	

Question Number	Acceptable Answer	Additional Guidance	Mark
3(b)(i)	<ul style="list-style-type: none"> calculates $\sum \Delta_f H(\text{products})$ 	(1) Example of calculation (+90.4 x 4) + (-241.8 x 6) = -1089.2	(3)
	<ul style="list-style-type: none"> $\sum \Delta_f H(\text{products}) - \Delta_r H$ 	(1) -1089.2 - (-904.8) = -184.4	
	<ul style="list-style-type: none"> calculates $\Delta_f H_{(NH_3)}$ for 1 mol ammonia 	(1) -184.4/4 = -46.1 (kJ mol ⁻¹) TE from M1 to M2 M3 can be awarded for an incorrect answer to M2 divided by 4 correct answer with no working scores 3 marks	

Question Number	Answer Acceptable	Additional Guidance	Mark
3(b)(ii)	<ul style="list-style-type: none"> <li data-bbox="367 309 685 336">• correct expression <li data-bbox="367 667 958 694">• correct evaluation of atom economy 	<p data-bbox="1189 272 1518 300"><u>Example of calculation</u></p> $\frac{4\text{NO}}{4\text{NO} + 6\text{H}_2\text{O}}$ <p data-bbox="1189 416 1234 443">OR</p> $\frac{4\text{NO}}{4\text{NH}_3 + 5\text{O}_2}$ <p data-bbox="1189 596 1653 624">may be shown as numbers only</p> $\frac{4(14 + 16)}{4(14 + 16) + 6(16 + 2)} \times 100$ <p data-bbox="1189 775 1234 802">OR</p> $\frac{4(14 + 16)}{4(14 + 3) + 5(16 \times 2)} \times 100$ <p data-bbox="1189 954 1890 1090"> = 53/52.6(316)(%) allow answer to 2 or 3 SF only correct answer with no working scores 2 marks 0.53/0.526 scores M1 only </p>	<p data-bbox="1944 233 2029 260">Mark</p> <p data-bbox="1957 1059 2011 1086">(2)</p>

Question Number	Acceptable Answer	Additional Guidance	Mark
3(c)(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> yield (of NO) decreases (1) increase in pressure shifts equilibrium (position) to the side of fewer moles (of gas molecules) (1) 	<p>if M1 and M2 are contradictory then do not award any marks</p> <p>allow 9 mol on LHS and 10 mol on RHS, may be shown above the equation</p> <p>allow more moles of product</p> <p>allow fewer moles of reactant</p> <p>allow marking points in either order</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
3(c)(ii)	<p>An answer that makes reference to the following points: (on increasing the pressure)</p> <ul style="list-style-type: none"> Rate increases because there are more molecules per unit volume (1) <p>so increase in frequency of collisions (between reacting molecules) (1)</p>	<p>allow increase in concentration of (gas) molecules</p> <p>allow any implication of more particles in a given volume, e.g. particles are closer together</p> <p>allow more collisions per unit time</p> <p>ignore just 'more collisions'/'more successful collisions' with no reference to time</p> <p>allow answers based on a solid catalyst</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
3(c)(iii)	<p>An answer that makes reference to:</p> <ul style="list-style-type: none"> heterogeneous: (the catalyst is in) a different phase/state to the reactants (1) increases the rate of the forward and backward / reverse reactions (1) 	ignore reference to products	(2)

Question Number	Acceptable Answer	Mark
3(d)	<p>The only correct answer is B</p> <p><i>A is not correct because there is no increase in number of particles</i></p> <p><i>C is not correct because distribution broadens as temperature rises, so peak is lower</i></p> <p><i>D is not correct because E_a is an intrinsic property of the reaction, not the applied temperature</i></p>	(1)

(Total for Question 3 = 14 marks)

Question Number	Acceptable Answer	Additional Guidance	Mark
4(a)(i)	Reagent • (concentrated) NaOH/KOH (1) Conditions • ethanol (solvent) <u>and</u> heat/warm (1)	do not award OH ⁻ or just 'hydroxide' do not award M1 if 'acidified' allow reflux M2 is dependent on M1 except for a near miss e.g. OH ⁻	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
4(a)(ii)	• Reagent: KCN/NaCN /potassium cyanide / sodium cyanide (1) • Reason: increases the number of carbon atoms in the carbon chain/ length of carbon chain (1)	ignore any mention of the solvent (aq ethanol) and conditions (reflux) do not award just CN ⁻ /cyanide/HCN	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
4(a)(iii)	An explanation that makes reference to the following: • heating increases rate (of reaction) (1) • no sealed tube would result in loss of ammonia (gas)/ reactants / gas (1)	ignore reference to activation energy/ starting the reaction/ reaction is endothermic ignore toxicity of reactants	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
4(a)(iv)	$\text{CH}_3\text{---CH}_2\text{---CH}_2\text{---OH}$	allow displayed/structural/skeletal formula ignore name do not award just C ₃ H ₇ OH	(1)


Question Number	Acceptable Answer	Mark
4(b)	<p>The only correct answer is B</p> <p><i>A is not correct because Z (3rd) is tertiary (fastest)</i></p> <p><i>C is not correct because Y (2nd) is primary (slower than X, secondary)</i></p> <p><i>D is not correct because X (1st) is secondary (slower than Z, tertiary)</i></p>	(1)

(Total for Question 4 = 8 marks)

Question Number	Acceptable Answer	Mark
5(a)(i)	<p>The only correct answer is B</p> <p><i>A is not correct because reaction is not substitution</i></p> <p><i>C is not correct because reaction is not substitution, nor nucleophilic</i></p> <p><i>D is not correct because reaction is not nucleophilic</i></p>	(1)

Question Number	Acceptable Answer	Mark
5(a)(ii)	<p>The only correct answer is C</p> <p><i>A is not correct because no C=C present</i></p> <p><i>B is not correct because no C=C present</i></p> <p><i>D is not correct because these are not stereoisomers</i></p>	(1)

Question	Acceptable Answer	Additional Guidance	Mark
----------	-------------------	---------------------	------

Number			
5(a)(iii)	<p>An answer which shows the following:</p> <ul style="list-style-type: none"> curly arrow from double bond to H atom of HCl/space between double bond and H atom of HCl (1) correct dipole on HCl molecule <u>and</u> curly arrow from H-Cl bond to Cl atom (1) intermediate with + charge shown on correct carbon (1) curly arrow from <u>lone pair</u> on chloride ion to correct carbon (1) 	<p><u>Example of mechanism</u></p>  <p>incorrectly drawn starting molecule loses M1, e.g. missing H or pentavalent carbon. incorrect starting molecule, e.g. butene will lose M3. if product is 1-chloropropane M3 only is lost. Other errors in end product lose M4 use of HBr in place of HCl loses M2 only use of H⁺ and Cl⁻ loses M1 and M2</p>	(4)

Question	Acceptable Answer	Additional Guidance	Mark
----------	-------------------	---------------------	------

Number			
5(b)(i)		<p>must show two repeat units fully displayed</p> <p>allow head to head, head to tail, tail to tail, syndiotactic and isotactic structures</p> <p>do not award any other type of formula</p> <p>ignore brackets and n</p>	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
5(b)(ii)	<p>An explanation that makes reference to the following:</p> <ul style="list-style-type: none"> (incineration produces) HCl/chlorinated molecules (1) which are corrosive/toxic /cause acid rain (1) 	<p>M2 is dependent on M1</p> <p>allow chlorine</p> <p>ignore carbon dioxide and its consequences</p> <p>allow adverse effect on ozone layer</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
5(b)(iii)	<p>An answer that makes reference to the following:</p> <p>any appropriate precautions to deal with toxic vapours/use fume cupboard etc.</p>	<p>allow good ventilation required</p> <p>allow gas mask/respirator</p> <p>do not award just mask</p> <p>ignore gloves, lab coat</p>	(1)
Question	Acceptable Answer	Additional Guidance	Mark

Number			
5(c)(i)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> at lower temperatures (below 50°C) the reaction will be slow (1) at higher temperatures (above 80°C) yield will be lower <u>because</u> (forward) reaction is exothermic (1) 	<p>allow reverse argument</p> <p>allow other products produced at higher temperatures</p>	(2)

Question Number	Acceptable Answer	Mark
5(c)(ii)	<p>The only correct answer is A</p> <p><i>B is not correct because separating funnel is inappropriate for an industrial process</i></p> <p><i>C is not correct because not a separation process</i></p> <p><i>D is not correct because both will react with alkaline solution</i></p>	(1)

(Total for Question 5 = 13 mark)

Question Number	Acceptable Answer	Additional Guidance	Mark
6(a)(i)	Reagent: <ul style="list-style-type: none"> B is hydrogen / H₂ (gas) (1) Condition: <ul style="list-style-type: none"> nickel/ Ni (catalyst) (1) 	mark independently allow any other suitable transition metal catalysts eg Pt, Pd ignore additional information relating to the support for the catalyst ignore references to heating/pressure/UV	(2)

Question Number	Acceptable Answer	Mark
6(a)(ii)	The only correct answer is C A is not correct because water is not involved B is not correct because there is no increase in number of oxygen atoms D is not correct because no substitution has taken place	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
6(a)(iii)	margarine	allow <u>liquid</u> coal allow butter substitute do not award just butter	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark																				
*6(b)	<p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning. Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning. The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="365 515 1189 786"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table border="1" data-bbox="365 895 1189 1358"> <thead> <tr> <th></th> <th>Number of marks awarded for structure and sustained lines of reasoning</th> </tr> </thead> <tbody> <tr> <td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.</td> <td>2</td> </tr> <tr> <td>Answer is partially structured with some linkages and lines of reasoning.</td> <td>1</td> </tr> <tr> <td>Answer has no linkages between points and is unstructured.</td> <td>0</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure and sustained lines of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2	Answer is partially structured with some linkages and lines of reasoning.	1	Answer has no linkages between points and is unstructured.	0	<p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning, scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> <p>In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.</p> <p>If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded do not deduct mark(s).</p> <p>Comment: Look for the indicative marking points first, then consider the mark for the structure of the answer and sustained line of reasoning.</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																						
6	4																						
5-4	3																						
3-2	2																						
1	1																						
0	0																						
	Number of marks awarded for structure and sustained lines of reasoning																						
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2																						
Answer is partially structured with some linkages and lines of reasoning.	1																						
Answer has no linkages between points and is unstructured.	0																						

	<p>Indicative content:</p> <ul style="list-style-type: none"> • calculate approximate mass of solute to be weighed out • details of how to weigh out required mass • transfer solute to beaker/conical flask <u>and</u> add distilled/deionised water <u>and</u> dissolve • transfer to (250 cm³) volumetric flask • add washings from beaker • make up to mark/line <u>and</u> shake/invert (to mix). 	<p>Ignore anything to do with oxidation even if incorrect</p> <p><u>example of calculation</u></p> $0.050 \text{ mol dm}^{-3} = 0.050 \times 118 \text{ g dm}^{-3}$ $= 5.90 \text{ g dm}^{-3}$ $= 1.47(5) \text{ g in } 250 \text{ cm}^3$ <p>do not award just 'weigh by difference'</p> <p>transfer of solute directly to volumetric flask gets IP3 and IP4 but must mention dissolving for IP3</p> <p>any mention of volumetric/graduated flask scores IP4</p> <p>direct transfer from weighing container to volumetric flask must mention washing of solute into the flask (e.g. through funnel).</p> <p>mix on its own is insufficient</p>	
--	--	--	--

(Total for Question 6 = 10 marks)

Question Number	Acceptable Answer	Additional Guidance	Mark
7(a)(i)	<ul style="list-style-type: none"> • ticks under titration numbers 2, 3, 4 (1) • 17.65 (cm³) (1) 	ignore X under Titration 1 <u>example of calculation</u> $\frac{17.60 + 17.70 + 17.65}{3} = 17.65$ scroll down as mean titre value may be written below (i) rather than in the table units not required must be 2 dp TE from M1 if Titration 1 has been ticked (17.74)	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
7(a)(ii)	<ul style="list-style-type: none"> • Phenolphthalein/ methyl orange (1) • colourless to pink / red to orange (1) 	M2 depends on M1 allow any indicator other than litmus or universal indicator allow minor errors in spelling of phenolphthalein but not phenyl... do not award red/pink-red for phenolphthalein nor yellow for methyl orange allow correct colour change for other indicators	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
7(a)(iii)	<ul style="list-style-type: none"> converts [acid] from g dm⁻³ to mol dm⁻³ (1) calculates moles of acid in 25 cm³ (1) calculates moles of sodium hydroxide in titre cm³ (1) converts moles of sodium hydroxide in titre to mol dm⁻³ and gives the answer 3 SF (1) 	<p><u>Example of calculation</u></p> <p>3.80/90.0 = *4.22 x 10⁻² (mol dm⁻³)</p> <p>ans to M1 x 25 x 10⁻³ 25 x 10⁻³ x *4.22 x 10⁻² = **1.0556 x 10⁻³ (mol) allow M1 and M2 in any order one mark only if not divided by 90.0</p> <p>ans to M2** x 2 = 1.0556 x 10⁻³ x 2 = ***2.111 x 10⁻³ (mol)</p> <p>= ans to M3*** x 1000/17.65 = 0.1196 = 0.120 (mol dm⁻³)</p> <p>correct answer with no working scores 4 marks</p>	(4)

Question Number	Acceptable Answer	Additional Guidance	Mark
7(b)(i)	<ul style="list-style-type: none"> burette uncertainty (1) pipette uncertainty (1) 	<p><u>Example of calculations</u></p> <p>0.05 x 2 x 100/ 17.65 = (±)0.567/0.57/0.6(%)</p> <p>0.06 x 100/25 = (±)0.24/0.2(%)</p> <p>ignore addition of the two uncertainties ignore SF</p>	(2)

Question Number	Acceptable Answer	Mark
7(b)(ii)	<p>The only correct answer is B</p> <p><i>A is not correct because the volume of NaOH needed is divided by 4, uncertainty is x4</i></p> <p><i>C is not correct because moles of acid is the same and uncertainty is the same.</i></p> <p><i>D is not correct because moles of acid halved and uncertainty doubled.</i></p>	<p>(1)</p>

(Total for Question 7 = 11 marks)

Question Number	Acceptable Answer	Additional Guidance	Mark
8(a)	$ \begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{H} \\ \quad \\ \text{H} \quad \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array} $	display all three methyl groups allow -OH do not award C-H-O	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
8(b)(i)	An answer that makes reference to one of the following: molecular ion/molecule fragments/is unstable		(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
8(b)(ii)	$ \begin{array}{c} \text{CH}_3-\overset{+}{\text{C}}-\text{CH}_3 \\ \\ \text{O}-\text{H} \end{array} $	allow + charge on any part of the ion/outside the structure but + must be shown allow displayed/structural/skeletal/ molecular formulae or any combination of these.	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
8(c)(i)	<ul style="list-style-type: none"> • calculation for bonds broken in the alcohol (*) (1) • calculation for bonds broken in oxygen <p style="text-align: center;">and</p> <p>total energy for bonds broken(**) (1)</p> <ul style="list-style-type: none"> • calculation for bonds made(***) (1) • calculation of $\Delta_c H$ (2-methylpropan-2-ol) with sign (1) 	<p><u>Example of calculation</u></p> $3(\text{C-C}) + 9(\text{C-H}) + (\text{C-O}) + (\text{O-H})$ $= (3 \times 347) + (9 \times 413) + 358 + 464 = (+)5580 \text{ (kJ mol}^{-1}\text{)}$ $6(\text{O=O}) = (6 \times 498) = (+)2988 \text{ (kJ mol}^{-1}\text{)}$ <p>total = + 5580 + 2988 = (+)8568 (kJ mol⁻¹) TE from ans * M1 + 2988</p> $= 8(\text{C=O}) + 10(\text{O-H})$ $= (8 \times 805) + (10 \times 464) = -11080 \text{ (kJ mol}^{-1}\text{)}$ $= +8568 - 11080 = -2512 \text{ (kJ mol}^{-1}\text{)}$ <p>allow TE for answer(**) + answer(***) units not required but if given they must be correct correct final answer with no working scores 4 marks</p>	(4)

Question Number	Acceptable Answer	Additional Guidance	Mark
8(c)(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> incomplete combustion (1) $\Delta_c H$ (2-methylpropan-2-ol) will be less negative /less exothermic than data book value (1) 	<p>mark independently</p> <p>do not award just lower/smaller/decreases/more positive</p> <p>allow reduce the magnitude (of the value)</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
8(c)(iii)	<p>An answer that makes reference to the following points:</p> <p>$\Delta_c H$ figures are at 298 K /data book bond energies refer to gaseous state</p> <p><u>and</u></p> <p>water and/or 2-methylpropan-2-ol are/is (both) liquid(s) (at 298 K)</p>	<p>allow just liquid involved</p> <p>do not award data book bond energies are mean (values)/not specific to 2-methylpropan-2-ol</p>	(1)

Question Number	Acceptable Answer	Mark
8(d)	<p>The only correct answer is D</p> <p><i>A is not correct because tertiary alcohol is not oxidised</i></p> <p><i>B is not correct because this is incorrect colour change for acidified dichromate</i></p> <p><i>C is not correct because this is incorrect colour change for these reagents</i></p>	(1)

(Total for Question 8 = 11 marks)

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom