
PHYSICS

9702/41

Paper 4 A Level Structured Questions

May/June 2018

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

IGCSE™ is a registered trademark.

This document consists of **14** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

| Question | Answer | Marks |
|------------|--|-----------|
| 1(a) | force proportional to product of masses and inversely proportional to square of separation | B1 |
| | idea of force between point masses | B1 |
| 1(b)(i) | velocity changes/direction of motion changes/there is an acceleration/there is a resultant force so not in equilibrium | B1 |
| 1(b)(ii)1. | gravitational force equals/is centripetal force | C1 |
| | $GMm/R^2 = mR\omega^2$ and $\omega = 2\pi/T$ or $Gm/R^2 = mv^2/R$ and $v = 2\pi r/T$ or $GMm/R^2 = mR(2\pi/T)^2$ | M1 |
| | convincing algebra leading to $k = GM/4\pi^2$ | A1 |
| 1(b)(ii)2. | correct use of R^3/T^2 for one planet (c gives 3.54×10^{21} ; e and g both give 3.56×10^{21}) | C1 |
| | $3.5(5) \times 10^{21} = (6.67 \times 10^{-11} \times M) / 4\pi^2$ $M = 2.1 \times 10^{33} \text{ kg}$ | A1 |
| | two or three values of R^3/T^2 correctly calculated and used in a valid way to find a value for M based on more than one k | B1 |

PUBLISHED

| Question | Answer | Marks |
|-----------------|--|--------------|
| 2(a)(i) | straight line through origin indicates acceleration \propto displacement | B1 |
| | negative gradient shows acceleration and displacement are in opposite directions | B1 |
| 2(a)(ii) | $a = -\omega^2 y$ and $\omega = 2\pi f$ | C1 |
| | $4.5 = (2\pi \times f)^2 \times 8.0 \times 10^{-3}$ (or other valid read-off) | |
| | $f = 3.8$ Hz | A1 |
| 2(b)(i) | maximum displacement upwards/above rest/above the equilibrium position | B1 |
| 2(b)(ii) | (just leaves plate when) acceleration = 9.81 ms^{-2} | C1 |
| | $9.81 = (2\pi \times 3.8)^2 \times y_0$ or $9.81 = 563 \times y_0$ | C1 |
| | amplitude = 17 mm | A1 |

PUBLISHED

| Question | Answer | Marks |
|-----------------|---|--------------|
| 3(a)(i) | sum of potential and kinetic energies (of molecules/atoms/particles) | B1 |
| | (energy of) molecules/atoms/particles in random motion | B1 |
| 3(a)(ii) | (in ideal gas) no intermolecular forces so no potential energy | B1 |
| | internal energy is (solely) kinetic energy (of particles) | B1 |
| | (mean) kinetic energy (of particles) proportional to (thermodynamic) temperature of gas | B1 |
| 3(b) | $pV = NkT$ | C1 |
| | $6.4 \times 10^6 \times 1.8 \times 10^4 \times 10^{-6} = N \times 1.38 \times 10^{-23} \times 298$ | C1 |
| | or | |
| | $pV = nRT$ and $N = n \times N_A$ | (C1) |
| | $6.4 \times 10^6 \times 1.8 \times 10^4 \times 10^{-6} = n \times 8.31 \times 298$ $n = 46.5$ (mol) $N = 46.5 \times 6.02 \times 10^{23}$ | (C1) |
| | $N = 2.8 \times 10^{25}$ | A1 |

PUBLISHED

| Question | Answer | Marks |
|-----------------|--|--------------|
| 4(a) | e.g. microphone weighing scales/pressure sensor lighters/spark generation watches/clocks/regulation of time | B1 |
| 4(b) | <u>pulses</u> (of ultrasound) | B1 |
| | reflected at boundaries (between media) | B1 |
| | (reflected pulses) detected by (ultrasound) generator | B1 |
| | Any three from: <ul style="list-style-type: none"> • time delay (between transmission and receipt) gives information about depth (of boundary) • intensity of reflected pulse gives information about (nature of) <u>boundary</u> • gel used to minimise reflection at skin/maximise transmission into skin • degree of reflection depends upon impedances of two media (at boundary) | B3 |

| Question | Answer | Marks |
|-----------|--|-------|
| 5(a)(i) | west to east | B1 |
| 5(a)(ii) | above the Equator | B1 |
| 5(a)(iii) | value in range $(1-300) \times 10^9$ Hz | A1 |
| 5(b)(i) | gain/dB = $10 \lg(P_2/P_1)$ | C1 |
| | $-195 = 10 \lg(P/3000)$ or $195 = 10 \lg(3000/P)$ | C1 |
| | power = 9.5×10^{-17} W | A1 |
| 5(b)(ii) | up-link has been (greatly) attenuated (before reaching satellite) or down-link signal must be (greatly) amplified (before transmission back to Earth) or up-link has (much) smaller intensity/power than down-link | B1 |
| | (different frequency) prevents down-link (signal) swamping up-link (signal) | B1 |

| Question | Answer | Marks |
|----------|---|-------|
| 6(a) | force per unit charge | B1 |
| 6(b) | $E = Q/(4\pi\epsilon_0 r^2)$ | C1 |
| | $2.0 \times 10^4 = Q / (4\pi \times 8.85 \times 10^{-12} \times 0.26^2)$ charge = 1.5×10^{-7} C | A1 |
| | | |
| 6(c) | charge (= $Q [52/26]^2$) = 4Q | C1 |
| | additional charge = 3Q | A1 |

PUBLISHED

| Question | Answer | Marks |
|-----------------|--|--------------|
| 7(a) | (capacitance =) charge / potential | M1 |
| | charge is (numerically equal to) charge on one plate | A1 |
| | potential is potential difference between plates | A1 |
| 7(b)(i) | $4.5 \times 10^{-6} \text{ C}$ | A1 |
| 7(b)(ii) | $9.0 \times 10^{-8} \text{ C}$ | A1 |
| 7(b)(iii) | capacitance = $(9.0 \times 10^{-8}) / 120$ | C1 |
| | $= 7.5 \times 10^{-10} \text{ F}$ | A1 |
| 7(c) | total capacitance is halved | B1 |
| | current is halved | B1 |

PUBLISHED

| Question | Answer | Marks |
|-----------------|---|--------------|
| 8(a)(i) | (fraction of) output is combined with the input | M1 |
| | output (fraction) subtracted/deducted from input | A1 |
| 8(a)(ii) | any two valid points e.g.: <ul style="list-style-type: none"> • greater bandwidth/gain constant over a larger range of frequencies/greater bandwidth • smaller gain | B2 |
| 8(b)(i) | gain = $(-)\frac{9600}{800}$ | C1 |
| | = -12 | A1 |
| 8(b)(ii) | 1. 1.2V | B1 |
| | 2. -6V | B1 |
| 8(b)(iii) | replace the 9600Ω resistor with an LDR | B1 |

PUBLISHED

| Question | Answer | Marks |
|----------|---|-----------|
| 9(a) | using Fleming's left-hand rule force on wire is upwards | B1 |
| | by Newton's third law, force on magnet is downwards | B1 |
| 9(b)(i) | $F = BIL$ | C1 |
| | $= 3.7 \times 10^{-3} \times 5.1 \times 8.5 \times 10^{-2}$ | A1 |
| | $= 1.6 \times 10^{-3} \text{ N}$ | |
| 9(b)(ii) | $F = 1.6 \times 10^{-3} \text{ N}$ | A1 |
| 9(c) | sketch: sinusoidal wave with two cycles | B1 |
| | amplitude $2.3 \times 10^{-3} \text{ N}$ | B1 |
| | period 0.05 s | B1 |

| Question | Answer | Marks |
|----------|---|-----------|
| 10(a) | induced <u>e.m.f.</u> proportional to rate | M1 |
| | of <u>change</u> of (magnetic) <u>flux</u> (linkage) or of <u>cutting</u> (magnetic) <u>flux</u> | A1 |
| 10(b) | current in coil produces flux | B1 |
| | (by Faraday's law) changing flux induces e.m.f. in ring | B1 |
| | current in ring causes field (around ring) | B1 |
| | (by Lenz's law) field around ring opposes field around coil | B1 |

PUBLISHED

| Question | Answer | Marks |
|-----------------|--|--------------|
| 11(a)(i) | packet/quantum/discrete amount of <u>energy</u> | M1 |
| | of electromagnetic radiation | A1 |
| 11(a)(ii) | (maximum) energy of emitted electrons is independent of intensity or no emission of electrons below the threshold frequency regardless of intensity or no emission of electrons when photon energy is less than work function (energy) regardless of intensity | B1 |
| 11(b) | in darkness: conduction band empty so high resistance | B1 |
| | in daylight: electrons in valence band absorb photons | B1 |
| | in daylight: electrons 'jump' to conduction band | B1 |
| | this leaves holes in valence band | B1 |
| | more charge carriers in daylight so resistance decreases | B1 |

PUBLISHED

| Question | Answer | Marks |
|-----------------|---|--------------|
| 12(a)(i) | $I = I_0 e^{-\mu x}$ | C1 |
| | $= I_0 \exp(-0.90 \times 2.8)$ | A1 |
| | $= 0.080 I_0$ | |
| 12(a)(ii) | $I = I_0 \exp[(-0.90 \times 1.5) \times (-3.0 \times 1.3)]$ | C1 |
| | $= I_0 (0.259 \times 0.20)$ | A1 |
| | $= 0.0052 I_0$ | |
| 12(b)(i) | difference in degrees of blackening | M1 |
| | between structures | A1 |
| 12(b)(ii) | large difference in intensities so good contrast | B1 |

| Question | Answer | Marks |
|-----------|---|-------------|
| 13(a) | emission of particles/radiation by unstable nucleus | B1 |
| | spontaneous emission | B1 |
| 13(b)(i) | use of graph to determine half-life = 14 minutes | B1 |
| | hence $\lambda = \ln 2 / (14 \times 60) \text{ (s}^{-1}\text{)}$ | C1 |
| | N at 14 minutes = 4.4×10^7 and $A = \lambda N$ | C1 |
| | activity = $4.4 \times 10^7 \times \ln 2 / (14 \times 60)$ = $3.6 \times 10^4 \text{ Bq}$ | A1 |
| | or | |
| | correct tangent drawn at time $t = 14$ minutes | (B1) |
| | magnitude of gradient of tangent identified as activity | (C1) |
| | correct working for gradient leading to activity | (C1) |
| | activity = $3.6 \times 10^4 \text{ Bq}$ | (A1) |
| 13(b)(ii) | $3.6 \times 10^4 = \lambda \times 4.4 \times 10^7$ or $\lambda = \ln 2 / (14.0 \times 60)$ | C1 |
| | $\lambda = 8.2 \times 10^{-4} \text{ s}^{-1}$ | A1 |