

**CAMBRIDGE**  
INTERNATIONAL EXAMINATIONS

**JUNE 2003**

**GCE A AND AS LEVEL**

**MARK SCHEME**

**MAXIMUM MARK: 60**

**SYLLABUS/COMPONENT: 9701/02**

**CHEMISTRY**  
**Theory 1 (Structured Questions)**

Page 1	Mark Scheme	Syllabus	Paper
	A/AS LEVEL EXAMINATIONS – JUNE 2003	9701	2

- 1 (a) Atoms which have the same number of protons (or same element) but different numbers of neutrons (1) [1]
- (b) (i)  $^{35}\text{Cl}$  (1)
- (ii)  $\text{H}^{37}\text{Cl}$  (1) [2]
- (c) H Cl line at 36 has rel. abundance of  $\left. \begin{array}{l} 90 \\ 38 \end{array} \right\}$   $\left. \begin{array}{l} 30 \\ 30 \end{array} \right\}$  (1)
- These show  $^{35}\text{Cl}$  and  $^{37}\text{Cl}$  in ratio 3:1 (1)  
[or use of 35 and 37] [2]
- (d) Mean of the two isotopes  $\frac{3 \times 35 + 1 \times 37}{4} = 35.5$  (1) [1]
- [Total: 6]**
- 2 (a) (i) That the volume of the gas molecules is negligible compared to the volume of gas (1)
- (ii) That there are no intermolecular forces  
OR collisions of the molecules are perfectly elastic  
Particles are in constant motion, losing no energy on collision (1) any two [2]
- (b)  $6.02 \times 10^{23}$  (1) [1]
- (c) (i)  $r = \underline{0.192}$  nm (1) Assume most candidates will work in  $\text{dm}^3$   
 $v = \frac{4}{3} \times 3.14 \times (1.92 \times 10^{-9})^3 = 2.96 \times 10^{-26} \text{ dm}^3 (2.96 \times 10^{-29} \text{ m}^3)$  (1)
- (ii)  $2.96 \times 10^{-26} \times \underline{6.02 \times 10^{23}}$  (1) =  $1.78 \times 10^{-2} \text{ dm}^3 (1.78 \times 10^{-5} \text{ m}^3)$  (1)
- (iii)  $24 \text{ dm}^3 (0.024 \text{ m}^3)$  (1)
- (iv)  $\frac{1.78 \times 10^{-2} \times 10^2}{24} = 0.074\%$  (1)
- (v) Some statement which connects with (a) (i) above (1) **max [5]**
- (d)
  - hot metals will react with oxygen in air (or nitrogen)
  - to form oxides/will burn out/to a powder
  - argon will not react
  - at high temperatures  $\text{O}_2$  and  $\text{N}_2$  in air will react to give  $\text{NO}_x$
NOT expansion of gases on heating any two [2]

**[Total: 10]**

Page 2	Mark Scheme	Syllabus	Paper
	A/AS LEVEL EXAMINATIONS – JUNE 2003	9701	2

3 (a)  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  (1) exothermic (1) [2]

(b) Pr. 50 atm upwards; Temp 400-600°C; catalyst of iron  
(1 each, conditions stated) [3]

(c) Too high a temp and equilibrium favours LHS, less ammonia at equilibrium (1)  
Too low a temp, rate too slow/not enough molecules have  $E_{act}$  (1) [2]

(d) (i) 
$$K_p = \frac{PNH_3^2}{PN_2 \times PH_2^3} \quad (1)$$

(ii) 
$$K_p = \frac{37.2^2}{44.8 \times 105.6^3} \quad (1)$$


$$= 2.62 \times 10^{-5} \text{ atm}^{-2} \quad (1) \text{ calculation and units} \quad [3]$$

(e) Excess (hence uncontrolled) nitrates leach out of fields into streams, seas (1)  
Bacteria or algae grow fast/use oxygen/clog up water (1)  
Balance destroyed/fish unable to live (1)  
Process called eutrophication (1) [3]

any 3


[Total: 13]

4 (a) (i)









[2]

(ii)




[2]

Page 3	Mark Scheme	Syllabus	Paper
	A/AS LEVEL EXAMINATIONS – JUNE 2003	9701	2



[Total: 9]



(b) Alkane (1) [1]

(c) (i) Not biodegradable/does not decompose/unreactive  
 Not affected by enzymes  
 Not attacked by aqueous or polar reagents found in tissues  
 Insoluble/does not absorb water/cotton absorbs water  
 NOT is stronger than cotton  
 [equivalent worthy points; they may overlap - but allow - max 2] [2]

Page 4	Mark Scheme	Syllabus	Paper
	A/AS LEVEL EXAMINATIONS – JUNE 2003	9701	2

- (ii) Alkanes react with oxygen (combustion)  
 Not possible in muscle (1)  
 also react with halogens/in U.V. light  
 muscle is internal and no halogens (1)  
 [ecf for alkene answers in (b)] [2]

[Total: 6]

- 6 (a)  $\frac{66.7}{12}$        $\frac{11.1}{1}$        $\frac{22.2}{16}$   
 = 5.5      = 11.1      = 1.3875  
 Divide by 1.3875  
 $C_4H_8O$  (1)       $48 + 8 + 16 = 72$       hence  $C_4H_8O$  (1) [2]

- (b) (i) orange ppt (1) red to yellow/crystals or solid  
 (ii) ketone (1)  
 (iii)  $CH_3CH_2COCH_3$  or butanone (1) [3]

- (c) (i)  $NaBH_4$       allow  $NaAlH_4$  ( $LiAlH_4$ ) (1)       $H_2/Ni$  or Pt  
 (ii) secondary alcohol (1)  
 (iii)  $CH_3CH_2CHOHCH_3$  (1)  
 [Allow ecf marks if (b) (iii) is butanal] [3]

[Total: 8]

- 7 (a) (i) e.g.  
 $CH_3CO_2C_3H_7$        $CH_3CO_2CH(CH_3)_2$        $CH_3CH_2CO_2C_2H_5$        $H-CO_2C_4H_9$   
 $C_3H_7CO_2CH_3$       + branches      any three [3]

- (ii)  $RCO_2R' + NaOH \rightarrow RCO_2Na$  (1) +  $R'OH$  (1)  
 $\rightarrow RCO_2H + R'OH$  (1) only [2]

- (b) (i) \* volatile, or liquids (1)      immiscible, with water (1)      smell (1)  
 and (ii)      any two [2]

- (c) (i) solvents, perfumes, flavourings, lotions, olive or palm oils      any two  
 and (ii) To make soap, to make Terylene      [2]  
 NOT polyesters

[Maximum Total: 8]