

**MARK SCHEME for the May/June 2012 question paper  
for the guidance of teachers**

**9702 PHYSICS**

**9702/42**

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2012 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9702	42

## Section A

- 1 (a) force proportional to product of masses and inversely proportional to square of separation (*do not allow square of distance/radius*)  
*either* point masses *or* separation @ size of masses
- M1  
A1 [2]
- (b) (i)  $\omega = 2\pi / (27.3 \times 24 \times 3600)$  *or*  $2\pi / (2.36 \times 10^6)$   
 $= 2.66 \times 10^{-6} \text{ rad s}^{-1}$
- M1  
A0 [1]
- (ii)  $GM = r^3 \omega^2$  *or*  $GM = v^2 r$   
 $M = (3.84 \times 10^5 \times 10^3)^3 \times (2.66 \times 10^{-6})^2 / (6.67 \times 10^{-11})$   
 $= 6.0 \times 10^{24} \text{ kg}$   
 (special case: uses  $g = GM/r^2$  with  $g = 9.81$ ,  $r = 6.4 \times 10^6$  scores max 1 mark)
- C1  
M1  
A0 [2]
- (c) (i) grav. force  $= (6.0 \times 10^{24}) \times (7.4 \times 10^{22}) \times (6.67 \times 10^{-11}) / (3.84 \times 10^8)^2$   
 $= 2.0 \times 10^{20} \text{ N}$  (*allow 1 SF*)
- C1  
A1 [2]
- (ii) *either*  $\Delta E_p = Fx$  because  $F$  constant as  $x \ll$  radius of orbit  
 $\Delta E_p = 2.0 \times 10^{20} \times 4.0 \times 10^{-2}$   
 $= 8.0 \times 10^{18} \text{ J}$  (*allow 1 SF*)
- B1  
C1  
A1 [3]
- or*  $\Delta E_p = GMm/r_1 - GMm/r_2$   
 Correct substitution  
 $8.0 \times 10^{18} \text{ J}$   
 ( $\Delta E_p = GMm/r_1 + GMm/r_2$  is incorrect physics so 0/3)
- C1  
B1  
A1
- 2 (a) energy  $= \frac{1}{2} m \omega^2 a^2$  and  $\omega = 2\pi f$   
 $= \frac{1}{2} \times 37 \times 10^{-3} \times (2\pi \times 3.5)^2 \times (2.8 \times 10^{-2})^2$   
 $= 7.0 \times 10^{-3} \text{ J}$   
 (allow  $2\pi \times 3.5$  shown as  $7\pi$ )
- C1  
M1  
A0 [2]
- Energy  $= \frac{1}{2} m v^2$  and  $v = r\omega$   
 Correct substitution  
 Energy  $= 7.0 \times 10^{-3} \text{ J}$
- (C1)  
(M1)  
(A0)
- (b)  $E_K = E_p$   
 $\frac{1}{2} m \omega^2 (a^2 - x^2) = \frac{1}{2} m \omega^2 x^2$  *or*  $E_K$  *or*  $E_p = 3.5 \text{ mJ}$   
 $x = a/\sqrt{2} = 2.8/\sqrt{2}$  *or*  $E_K = \frac{1}{2} m \omega^2 (a^2 - x^2)$  *or*  $E_p = \frac{1}{2} m \omega^2 x^2$   
 $= 2.0 \text{ cm}$   
 ( $E_K$  *or*  $E_p = 7.0 \text{ mJ}$  scores 0/3)
- C1  
C1  
A1 [3]
- Allow:  $k = 17.9$   
 $E = \frac{1}{2} kx^2$   
 $x = 2.0 \text{ cm}$
- (C1)  
(C1)  
(A1)

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9702	42

	(c) (i) graph: horizontal line, y-intercept = 7.0 mJ with end-points of line at +2.8 cm and –2.8 cm	B1	[1]
	(ii) graph: reasonable curve with maximum at (0,7.0) end-points of line at (–2.8, 0) and (+2.8, 0)	B1 B1	[2]
	(iii) graph: inverted version of (ii) with intersections at (–2.0, 3.5) and (+2.0, 3.5) (Allow marks in (iii), but not in (ii), if graphs K & P are not labelled)	M1 A1	[2]
	(d) <u>gravitational potential</u> energy	B1	[1]
3	(a) sum of potential energy and kinetic energy of atoms/molecules/particles reference to random (distribution)	M1 A1	[2]
	(b) (i) as lattice structure is 'broken'/bonds broken/forces between molecules reduced (not molecules separate) no change in kinetic energy, potential energy increases internal energy increases	B1 M1 A1	[3]
	(ii) <i>either</i> molecules/atoms/particles move faster/ $\langle c^2 \rangle$ is increasing <i>or</i> kinetic energy increases with temperature (increases) no change in potential energy, kinetic energy increases internal energy increases	B1 M1 A1	[3]
4	(a) (i) as $r$ decreases, energy decreases/work got out (due to <u>attraction</u> ) so point mass is negatively charged	M1 A1	[2]
	(ii) electric potential energy = charge $\times$ electric potential electric field strength is potential gradient field strength = gradient of potential energy graph/charge	B1 B1 A0	[2]
	(b) tangent drawn at (4.0, 14.5) gradient = $3.6 \times 10^{-24}$ (for $\langle \pm 0.3$ allow 2 marks, for $\langle \pm 0.6$ allow 1 mark) field strength = $(3.6 \times 10^{-24}) / (1.6 \times 10^{-19})$ = $2.3 \times 10^{-5} \text{ V m}^{-1}$ (allow ecf from gradient value) (one point solution for gradient leading to $2.3 \times 10^{-5} \text{ V m}^{-1}$ scores 1 mark only)	B1 A2 A1	[4]

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9702	42

- 5 (a) (long) straight conductor carrying current of 1 A  
current/wire normal to magnetic field  
(for flux density 1 T,) force per unit length is  $1 \text{ N m}^{-1}$
- (b) (i) (originally) downward force on magnet (due to current)  
by Newton's third law (allow "N3")  
upward force on wire
- (ii)  $F = BIL$   
 $2.4 \times 10^{-3} \times 9.8 = B \times 5.6 \times 6.4 \times 10^{-2}$   
 $B = 0.066 \text{ T}$  (need 2 SF)  
(g missing scores 0/2, but g = 10 leading to 0.067 T scores 1/2)
- (c) new reading is  $2.4\sqrt{2} \text{ g}$   
either changes between +3.4 g and -3.4 g  
or total change is 6.8 g
- 6 (a) oil drop charged by friction/beta source  
between parallel metal plates  
plates are horizontal  
adjustable potential difference/field between plates  
until oil drop is stationary  
 $mg = q \times V/d$   
symbols explained  
oil drop viewed through microscope  
m determined from terminal speed of drop (when p.d. is zero)  
(any two extras, 1 each)
- (b)  $3.2 \times 10^{-19} \text{ C}$
- 7 (a) minimum energy to remove an electron from the metal/surface
- (b) gradient =  $4.17 \times 10^{-15}$  (allow 4.1 → 4.3)  
 $h = 4.15 \times 10^{-15} \times 1.6 \times 10^{-19}$  or  $h = 4.1 \text{ to } 4.3 \times 10^{-15} \text{ eVs}$   
 $= 6.6 \times 10^{-34} \text{ Js}$
- (c) graph: straight line parallel to given line  
with intercept at any higher frequency  
intercept at between  $6.9 \times 10^{14} \text{ Hz}$  and  $7.1 \times 10^{14} \text{ Hz}$

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9702	42

- 8 (a) nuclei having same number of protons/proton (atomic) number  
different numbers of neutrons/neutron number  
(allow second mark for nucleons/nucleon number/mass number/atomic  
mass if made clear that same number of protons/proton number)
- B1  
B1 [2]
- (b) probability of decay per unit time is the decay constant  
 $\lambda = \ln 2 / t_{1/2}$   
 $= 0.693 / (52 \times 24 \times 3600)$   
 $= 1.54 \times 10^{-7} \text{ s}^{-1}$
- C1  
C1  
A1 [3]
- (c) (i)  $A = A_0 \exp(-\lambda t)$   
 $7.4 \times 10^6 = A_0 \exp(-1.54 \times 10^{-7} \times 21 \times 24 \times 3600)$   
 $A_0 = 9.8 \times 10^6 \text{ Bq}$   
(alternative method uses 21 days as 0.404 half-lives)
- C1  
A1 [2]
- (ii)  $A = \lambda N$  and  $\text{mass} = N \times 89 / N_A$   
 $\text{mass} = (9.8 \times 10^6 \times 89) / (1.54 \times 10^{-7} \times 6.02 \times 10^{23})$   
 $= 9.4 \times 10^{-9} \text{ g}$
- C1  
A1 [2]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9702	42

## Section B

- 9 (a)** e.g. infinite input impedance/resistance  
zero output impedance/resistance  
infinite (open loop) gain  
infinite bandwidth  
infinite slew rate  
*(any four, one mark each)* B4 [4]
- (b)** graph: square wave M1  
180° phase change A1  
amplitude 5.0 V A1 [3]
- (c)** correct symbol for LED M1  
diodes connected correctly between  $V_{OUT}$  and earth A1  
diodes identified correctly A1 [3]  
*(special case: if diode symbol, not LED symbol, allow 2<sup>nd</sup> and 3<sup>rd</sup> marks to be scored)*
- 10 (a)** e.g. beam is divergent/obeys inverse square law  
absorption (in block)  
scattering (of beam in block)  
reflection (at boundaries)  
*(any two sensible suggestions, 1 each)* B2 [2]
- (b) (i)**  $I = I_0 \exp(-\mu x)$  C1  
 $I_0/I = \exp(0.27 \times 2.4)$   
= 1.9 A1 [2]
- (ii)**  $I_0/I = \exp(0.27 \times 1.3) \times \exp(3.0 \times 1.1)$  C1  
= 1.42 × 27.1  
= 38.5 A1 [2]
- (c)** *either* much greater absorption in bone than in soft tissue  
*or*  $I_0/I$  much greater for bone than soft tissue B1 [1]
- 11 (a) (i)** loss of (signal) power B1 [1]
- (ii)** unwanted power (on signal)  
that is random M1  
A1 [2]
- (b)** for digital, only the 'high' and the 'low' / 1 and 0 are necessary M1  
variation between 'highs' and 'lows' caused by noise not required A1 [2]
- (c)** attenuation =  $10 \lg(P_2 / P_1)$  C1  
*either*  $195 = 10 \lg\{2.4 \times 10^3 / P\}$   
*or*  $-195 = 10 \lg(P / 2.4 \times 10^3)$  C1  
 $P = 7.6 \times 10^{-17} \text{ W}$  A1 [3]

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9702	42

- 12 (a) (i) modulator B1 [1]
- (ii) serial-to-parallel converter (*accept series-to-parallel converter*) B1 [1]
- (b) (i) enables one aerial to be used for transmission and receipt of signals A1 [1]
- (ii) all bits for one number arrive at one time B1  
bits are sent out one after another B1 [2]