

Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel
Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Friday 22 May 2020

Afternoon (Time: 1 hour 30 minutes)

Paper Reference **9FM0/3A**

Further Mathematics

Advanced

Paper 3A: Further Pure Mathematics 1

You must have:

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P62673A

©2020 Pearson Education Ltd.

1/1/1/1/1/1/



Pearson

1. Use l'Hospital's Rule to show that

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{(e^{\sin x} - \cos(3x) - e)}{\tan(2x)} = -\frac{3}{2}$$

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2.




Figure 1

Figure 1 shows a sketch of the vertical cross-section of the entrance to a tunnel. The width at the base of the tunnel entrance is 2 metres and its maximum height is 3 metres.

The shape of the cross-section can be modelled by the curve with equation $y = f(x)$ where

$$f(x) = 3 \cos\left(\frac{\pi}{2}x^2\right) \quad x \in [-1, 1]$$

A wooden door of uniform thickness 85 mm is to be made to seal the tunnel entrance.

Use Simpson's rule with 6 intervals to estimate the volume of wood required for this door, giving your answer in m^3 to 4 significant figures.

(6)



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area consisting of 24 horizontal lines.

(Total for Question 2 is 6 marks)



3. The points A , B and C , with position vectors $\mathbf{a} = 3\mathbf{i} - 2\mathbf{j} + \mathbf{k}$, $\mathbf{b} = \mathbf{i} + 4\mathbf{j} + 5\mathbf{k}$ and $\mathbf{c} = -2\mathbf{i} + 3\mathbf{j} + 3\mathbf{k}$ respectively, lie on the plane Π

(a) Find $\vec{AB} \times \vec{AC}$ (3)

(b) Find an equation for Π in the form $\mathbf{r} \cdot \mathbf{n} = p$ (2)

The point D has position vector $8\mathbf{i} + 7\mathbf{j} + 5\mathbf{k}$

(c) Determine the volume of the tetrahedron $ABCD$ (4)

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA



Question 3 continued

Lined area for writing the answer to Question 3.

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA



4.

$$f(x) = x^4 \sin(2x)$$

Use Leibnitz's theorem to show that the coefficient of $(x - \pi)^8$ in the Taylor series expansion of $f(x)$ about π is

$$\frac{a\pi + b\pi^3}{315}$$

where a and b are integers to be determined.

(8)

$$\left[\begin{array}{l} \text{The Taylor series expansion of } f(x) \text{ about } x = k \text{ is given by} \\ f(x) = f(k) + (x - k)f'(k) + \frac{(x - k)^2}{2!}f''(k) + \dots + \frac{(x - k)^r}{r!}f^{(r)}(k) + \dots \end{array} \right]$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



6. A physics student is studying the movement of particles in an electric field. In one experiment, the distances in micrometres of two moving particles, A and B , from a fixed point O are modelled by

$$d_A = |5t - 31|$$

$$d_B = |3t^2 - 25t + 8|$$

respectively, where t is the time in seconds after motion begins.

- (a) Use algebra to find the range of time for which particle A is further away from O than particle B is from O .

(8)

It was recorded that the distance of particle B from O was less than the distance of particle A from O for approximately 4 seconds.

- (b) Use this information to assess the validity of the model.

(2)



Question 6 continued

Lined area for writing the answer to Question 6. The page contains 30 horizontal lines.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



7. The points $P(9p^2, 18p)$ and $Q(9q^2, 18q)$, $p \neq q$, lie on the parabola C with equation

$$y^2 = 36x$$

The line l passes through the points P and Q

(a) Show that an equation for the line l is

$$(p + q)y = 2(x + 9pq) \quad (3)$$

The normal to C at P and the normal to C at Q meet at the point A .

(b) Show that the coordinates of A are

$$(9(p^2 + q^2 + pq + 2), -9pq(p + q)) \quad (7)$$

Given that the points P and Q vary such that l always passes through the point $(12, 0)$

(c) find, in the form $y^2 = f(x)$, an equation for the locus of A , giving $f(x)$ in simplest form.

(4)



DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued



8.
$$f(x) = \frac{3}{13 + 6\sin x - 5\cos x}$$

Using the substitution $t = \tan\left(\frac{x}{2}\right)$

(a) show that $f(x)$ can be written in the form

$$\frac{3(1+t^2)}{2(3t+1)^2+6} \quad (3)$$

(b) Hence solve, for $0 < x < 2\pi$, the equation

$$f(x) = \frac{3}{7}$$

giving your answers to 2 decimal places where appropriate.

(5)

(c) Use the result of part (a) to show that

$$\int_{\frac{\pi}{3}}^{\frac{4\pi}{3}} f(x) dx = K \left(\arctan\left(\frac{\sqrt{3}-9}{3}\right) - \arctan\left(\frac{\sqrt{3}+3}{3}\right) + \pi \right)$$

where K is a constant to be determined.

(8)



Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

Area for writing answers to Question 8, consisting of multiple horizontal lines.

(Total for Question 8 is 16 marks)

TOTAL FOR PAPER IS 75 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

