MARK SCHEME for the May/June 2013 series

0625 PHYSICS

0625/31

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0625	31

NOTES ABOUT MARK SCHEME SYMBOLS & OTHER MATTERS

- M marks are method marks upon which further marks depend. For an M mark to be scored, the point to which it refers **must** be seen in a candidate's answer. If a candidate fails to score a particular M mark, then none of the dependent marks can be scored.
- B marks are independent marks, which do not depend on other marks. For a B mark to be scored, the point to which it refers must be seen specifically in the candidate's answers.
- A marks In general A marks are awarded for final answers to numerical questions.
 If a final numerical answer, eligible for A marks, is correct, with the correct unit and an acceptable number of significant figures, all the marks for that question are normally awarded.
 It is very occasionally possible to arrive at a correct answer by an entirely wrong approach. In these rare circumstances, do not award the A marks, but award C marks on their merits. However, correct numerical answers with no working shown gain all the marks available.
- C marks are compensatory marks in general applicable to numerical questions. These can be scored even if the point to which they refer are not written down by the candidate, **provided subsequent working gives evidence that they must have known it.** For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct substitution or working which shows he knew the equation, then the C mark is scored. A C mark is not awarded if a candidate makes two points which contradict each other. Points which are wrong but irrelevant are ignored.
- brackets () around words or units in the mark scheme are intended to indicate wording used to clarify the mark scheme, but the marks do not depend on seeing the words or units in brackets, e.g. 10 (J) means that the mark is scored for 10, regardless of the unit given.
- <u>underlining</u> indicates that this <u>must</u> be seen in the answer offered, or something very similar.
- OR / or indicates alternative answers, any one of which is satisfactory for scoring the marks.
- e.e.o.o. means 'each error or omission'.
- o.w.t.t.e. means 'or words to that effect'.
- Spelling Be generous about spelling and use of English. If an answer can be understood to mean what we want, give credit. However, beware of and do not allow ambiguities, accidental or deliberate: e.g. spelling which suggests confusion between reflection / refraction / diffraction / thermistor / transistor / transformer.
- Not/NOT Indicates that an incorrect answer is not to be disregarded, but cancels another otherwise correct alternative offered by the candidate i.e. right plus wrong penalty applies.
- Ignore Indicates that something which is not correct or irrelevant is to be disregarded and does not cause a right plus wrong penalty.

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0625	31

e.c.f. meaning 'error carried forward' is mainly applicable to numerical questions, but may in particular circumstances be applied in non-numerical questions. This indicates that if a candidate has made an earlier mistake and has carried an incorrect value forward to subsequent stages of working, marks indicated by ecf may be awarded, provided the subsequent working is correct, bearing in mind the earlier mistake. This prevents a candidate being penalised more than once for a particular mistake, but **only** applies to marks annotated e.c.f.

Significant Figures

Answers are normally acceptable to any number of significant figures \dot{u} 2. Accept answers that round to give the correct answer to 2 s.f. Any exceptions to this general rule will be specified in the mark scheme.

Units Deduct one mark for each incorrect or missing unit from a final answer that would otherwise gain all the marks available for that answer: maximum 1 per question.

Arithmetic errors

Deduct one mark if the **only** error in arriving at a final answer is clearly an arithmetic one.

Transcription errors

Deduct one mark if the only error in arriving at a final answer is because given or previously calculated data has clearly been misread but used correctly.

Fractions e.g. ¹/₂, ¹/₄, 1/10 etc. are only acceptable where specified.

	Page 4					Mark Sc	heme			Syllabu	JS	Paper	
					IGCS	E – May	/June 2	013		0625		31	
1	(a)	 (density =) mass/volume OR mass per unit volume OR m/V with symbols explained 								E	31		
	(b)	(i)	(vol = 22	=) mass/c .48 cm³	ensity O to 2 or n	R 60.7/2 nore sig.	.70 figs						C1 41
		(ii)	OR 2	A × (avera 22.48 / (50 1499 cm) × 30)								C1 41
	(c)	(i)	micr	ometer/sc	rew gau	ge / (verr	nier/digit	al) callipers	5			E	31
		(ii)		ck zero of ce / fold s		sed / cut	sheet ir	ito several	pieces	/ detail of	how to use		31
			OR r calcu		hickness	of sever	al piece	<u>places</u> s together divide ansv	ver by	number of	measuren	nents/	31 31
			proo	oo, piacoo								[Total	
2	(a)			e or circle e or circle									31 31
	(b)	(i)	4.07	– 4.1 (s)								E	31
		(ii)	ÔR ($u)/t$ OR Δ other correction of the correctio	ect value	s from gi	raph	f 40 ÷ (ans s/s	s. to (b))(i))			C1 41
		(iii)	area OR s 82 m	$s = ut + \frac{1}{2}$	aph OR ½ at ² OR v	$\frac{1}{2}(u+v)^{2}$ = u^{2} + 2	t OR ½ ∷ ?as OR r	× 40 × (ans numbers su	. to (b) Ibstitute	(i)) ed			C1 A1
	(c)	gra	ph co	ntinues in	straight	line to 6	S					[Total	∃1 8]

	Page 5			Mark Scheme	Syllabus	Paper			
				IGCSE – May/June 2013	0625	31			
3	(a)	(i)		(loss of P.E. =) mgh OR 92 × 10 × 1500 1.38 × 10 ⁶ J correct use of mgh with h = 500 or 2000 gains 1 ma	irk only	C1 A1			
		. ,		(K.E. =) $\frac{1}{2} mv^2$ OR $\frac{1}{2} \times 92 \times 52^2$ 1.244 × 10 ⁵ J at least 2 sig. figs		C1 A1			
	(a)	 (ii) difference is due to: (work done in overcoming) air resistance/drag OR energy converted to/lost as heat (by air resistance/drag) 							
	(b)	(i)	incre	eases		B1			
		(ii)	920	Ν		B1			
						[Total 7]			
4	(a)	(i)		ntion of vacuum OR glass is a poor conductor vacuum/gap between walls has no molecules/atoms	s/particles	B1			
		 (ii) surface/silver (of walls) is good reflector/poor absorber (of radiation) surface/silver (of walls) is poor emitter (of radiation) 							
	(b)			opper/lid/bung/cover/top to reduce/prevent (loss of h on/radiation/evaporation OR to prevent steam/hot va		M1 B1			
			made of insulator OR example of insulator to reduce/prevent (loss of heat by) convection/radiation/evaporation OR to prevent steam/hot air leaving						
						[Total 6]			
5	(a)	(i) a (i)	•	ii) marked together to maximum of 3 marks ecules escape/leave the liquid/form gas or vapour		B1			
		(ii)		poration OR heat/(thermal) energy needed for evapo (er) molecules/high(er) energy molecules escape	pration leaves sweat	cooler B1			
		OR slow(er) molecules left behind heat flows from body to warm the sweat (so body cools)							
	(b)	(i)		=) <i>mc∆θ</i> OR <i>mcT</i> OR 60 × 4000 × 0.50 × 10 ⁵ J / 120 kJ		C1 A1			
		(ii) $Q = mL$ in any form OR (m =) Q/L OR either with numbers $(m = 1.2 \times 10^5 / 2.4 \times 10^6 =) 0.05$ kg e.c.f from (b)(i)							
						[Total 7]			

	Page 6		i	Mark Scheme	Syllabus	Paper			
				IGCSE – May/June 2013	0625	31			
6	(a)	 (i) (pressure =) force/area OR force per unit area OR (P =) F/A with symbols explained 							
		 (ii) molecules collide with/hit walls/surface (of box) molecule(s) exert force on wall pressure is total force / force of all molecules divided by (total) area of wall 							
	(b)	(i) $(P =) h\rho g \text{ OR in words OR } 0.25 \times 13600 \times 10$ 34 000 Pa OR N/m ² allow 1 mark for <i>h</i> = 250 used and 3.4×10^7 Pa obtained							
		 (<i>P</i> = 1.02 × 10⁵ − 34 000) 68 000 Pa or N/m² e.c.f. from (b)(i) only if (b)(i) is less than 1.02 × 10⁵ 							
						[Total 7]			
7	(a)	two of: ray <u>through</u> centre of lens undeviated ray parallel to axis refracted to right hand focus rays through left hand focus refracted parallel to axis							
		rays extrapolated to a point							
		accuracy marks: image 6 cm from lens image 6 cm high							
	(b)) image is virtual/not real <u>AND</u> cannot be seen on screen OR no rays come from (position of) image							

	Page 7			Mark Scheme	Syllabus	Paper	
				IGCSE – May/June 2013	0625	31	
8	(a)	15–	25 H	z to 15 000–25000 Hz / 15–25 kHz		B1	
	clos			ion) where air layers/molecules/particles are pushec er (than normal) (region) where (air) pressure raised/air (more) comp	-	-	
	((ii)	· •	ion) where air layers/molecules are pushed apart/fa (region) where (air) pressure reduced/air expanded	r(ther) apart (than	normal) B1	
	(c)	(c) (i) (sound is) loud(er) OR volume (of sound is) increased					
	((ii)	sour	nd has a higher frequency/pitch OR higher note (hea	ard)	B1	
				0 OR 1.6 (s) seen OR <i>v</i> = 2 <i>d</i> /1.9 OR 500 (m) seen OR <i>v</i> = (2 <i>d</i> + 500)/3.5		C1 C1	
				= 500 / 1.6 =) 312.5 m/s at least 2 sig. figs		A1	
						[Total 8]	
9	(a)	(i)	all la	amps off			
	((ii)	12 Ω	2 lamps (only) on		B1	
	(iii) 4 <u>(</u>		4 Ω	lamps (only) on			
	(b)	(i)	12 V	/		B1	
	((ii)		// <i>R</i> in any form OR <i>V/R</i> OR 12/12		C1	
				A OR 1 A . from (b)(i)		A1	
				n 4 Ω lamp = 3 (A) (current in 12 Ω lamp is in (b)(ii))		C1 C1	
		(<i>P</i> =) <i>IV</i> OR <i>I</i> ² <i>R</i> (<i>P</i> =) 36 W for 4 Ω lamp; <i>P</i> = 12 W for 12 Ω lamp e.c.f. from (b)(ii)					
		OR					
		(P = (P =	(C1) (C1)				
		(<i>P</i> = OR	(A1)				
		(P =	(B1)				
		San 4 Ω	(M1) (A1)				
						[Total 7]	

	Pa	ige 8		Mark Scheme	Syllabus	Paper
				IGCSE – May/June 2013	0625	31
10	(a)	arro	ows cl	concentric circles centred on wire lockwise on each circle / at least one circle of circles increasing as radius increases		B1 B1 B1
	(b)	(i)	arrov	w pointing down on side AB, up on side CD		B1
		(ii)	line (OR h	es on AB and CD are opposite OR up and down an (so cause rotation) nave moments in same sense / direction cause couple / torque	d separated / not ii	n same B1
		(iii)	OR k vertio	verse current in loop or keep current in AB or CD ir keep current on side near a pole in the same direct cal every half turn		
			so th rotati OR s	ion continues (in same direction) so that rotation doesn't reverse its direction		B1
				to maintain sense/direction of moments/couple coil turns more than half a revolution		B1
						[Total 7]
11	(a)	(i)	•	otons utrons		B1 B1
		(ii)	a (fa	st moving) electron		B1
	(b) ele		ctron/e	electrons removed from/gained by the molecule		B1
	(c)	(i)	OR t	e because particle is charged the force on the particles is perpendicular to their pa direction of force changes as direction of motion ch		B1
		(ii)	α-ра	article <u>curve</u> up the page in at least half of width of t	field	B1
			curva	article <u>curve</u> opposite to α -particle curve OR down pature anywhere ller radius of β path clear	bage if α line has n	o B1 B1
			Sindi			
						[Total 8]