

**MARK SCHEME for the October/November 2010 question paper
for the guidance of teachers**

9700 BIOLOGY

9700/36

Paper 32 (Advanced Practical Skills 2),
maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2010	9700	36

Question	Expected Answers	Additional guidance
1 (a) (i)	Decide on the temperatures you plan to use in the range (between) 25°C to 45°C. Record the temperatures you have chosen in the space below.	[2]
MMO decisions 2	[1] at least 5 temperatures;	
	[1] one temp. 25°C to 29°C AND one temp 40°C to 45°C AND any three with two even intervals 3 or more degrees;	
(ii) Prepare the space below and record your results.		[4]
PDO recording 2	[1] Reject • if any units in body of table • only t	Must have units
	table with all cells drawn AND heading (top or left) temperature °C;	
	[1] Reject • if units in body of table • if headings for volumes or stages (heading) time with units;	
MMO collection 2	[1] temperatures recorded highest to lowest AND first set of times recorded in whole seconds;	
	[1] time at the lowest temperature is greater than the next temperature;	Allow • only if 3 or more results
(iii) From your results, state the temperature at which the activity of the enzyme is lowest.		[1]
ACE interpretation 1	[1] temperature with longest time AND with units, °C;	

(iv) Identify <i>two</i> significant sources of error in this investigation.			[2]
ACE interpretation max 2		cause of error	error
	[1]	(dependent) stage 3 or end-point clots stick small clots coagulation milk drains back slowly	idea of seeing determining judging when;
	[1]	(standardised variables) rotation or angle;	AND idea of not constant/different not same timing delayed;
	[1]	shaking or mixing or E/enzyme starts to react;	
	[1]	E/enzyme temperature; (as milk)/AW	
	[1]	(independent variable) temperature or test-tube removed from water-bath	idea of not constant/not maintained decreasing cools down;
			Max 2
(v) Describe a suitable control for this investigation.			[1]
Reject if give two.			
ACE improvement 1	[1]	boil enzyme;	

(vi) Suggest how you could make this investigation as reliable as possible.			[1]																																																
ACE improvements MAX 1	C control of any relevant variable [1]	equilibrate milk and enzyme to temp. separately then mix Or use thermostatically controlled water bath Or keep tube in water bath during rotation;																																																	
	R1 improve method to get repeat data [1]	repeat	AND calculate or find mean/average;																																																
			max 1																																																
(b) (i) Three of the values in Table 1.1 are anomalous. Draw a circle around each of these values.			[1]																																																
(ii) Complete the Table 1.1 by calculating the missing value.			[1]																																																
MMO decisions 1 ACE interpretation 1	[1]	circles around <u>8.2, 4.9, 1.1</u> ;																																																	
	[1]	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th colspan="7">activity of milk clotting enzyme / arbitrary units</th> </tr> <tr> <th>pH of milk</th> <th>trial 1</th> <th>trial 2</th> <th>trial 3</th> <th>trial 4</th> <th>trial 5</th> <th>mean</th> </tr> </thead> <tbody> <tr> <td>6.02</td> <td>8.8</td> <td>8.7</td> <td>8.9</td> <td>8.2</td> <td>8.7</td> <td>8.8</td> </tr> <tr> <td>6.22</td> <td>6.8</td> <td>6.8</td> <td>6.8</td> <td>6.7</td> <td>6.9</td> <td>6.8</td> </tr> <tr> <td>6.40</td> <td>4.9</td> <td>4.3</td> <td>4.4</td> <td>4.3</td> <td>4.4</td> <td>4.4</td> </tr> <tr> <td>6.64</td> <td>1.1</td> <td>1.0</td> <td>1.0</td> <td>0.9</td> <td>1.0</td> <td>1.0</td> </tr> <tr> <td>6.70</td> <td>0.7</td> <td>0.6</td> <td>1.1</td> <td>0.5</td> <td>0.7</td> <td>0.6</td> </tr> </tbody> </table>		activity of milk clotting enzyme / arbitrary units							pH of milk	trial 1	trial 2	trial 3	trial 4	trial 5	mean	6.02	8.8	8.7	8.9	8.2	8.7	8.8	6.22	6.8	6.8	6.8	6.7	6.9	6.8	6.40	4.9	4.3	4.4	4.3	4.4	4.4	6.64	1.1	1.0	1.0	0.9	1.0	1.0	6.70	0.7	0.6	1.1	0.5	0.7
activity of milk clotting enzyme / arbitrary units																																																			
pH of milk	trial 1	trial 2	trial 3	trial 4	trial 5	mean																																													
6.02	8.8	8.7	8.9	8.2	8.7	8.8																																													
6.22	6.8	6.8	6.8	6.7	6.9	6.8																																													
6.40	4.9	4.3	4.4	4.3	4.4	4.4																																													
6.64	1.1	1.0	1.0	0.9	1.0	1.0																																													
6.70	0.7	0.6	1.1	0.5	0.7	0.6																																													
		8.8 Allow 8.7																																																	

(iii) Plot a graph of the data shown in Table 1.1. [4]				
PDO layout 4	O [1]	x-axis pH	AND y-axis activity (/) arbitrary units or au;	Must have units
	S [1]	Reject if awkward scale scale as 0.2 to 2 cm Origin must be labelled as 6 or 6.02	AND 2 to 2 cm;	error carried forward if incorrect O then scale x-axis 2 to 2 cm and y-axis 0.2 to 2 cm. must use more than half grid in x and y.
	P [1]	Reject plotting if scale is awkward if only dots/blobs or blobs in circles correct plotting using crosses/dots in circle only;	intersection of cross must be clear to show plot.	
	L [1]	straight line through points; error carried forward if scale or plotting incorrect 6.02 8.8 or 8.7 or ecf 6.22 6.8 6.40 4.4 6.64 1.0 6.70 0.6	quality – not thick, not feathery for the complete line. joining plots – <ul style="list-style-type: none"> • <u>ruled lines plot to plot</u> • <u>line of best fit</u> • <u>curve through all plots</u> 	

(iv) Explain the relationship between pH and the enzyme shown in the data.			[3]
ACE conclusions 3	[1]	(in correct context of pH and effect on activity) structure of protein or substrate or enzyme or active site or bonds	changed/altered/destroyed/no longer complementary broken;
	[1]	(in correct context of increase in pH and decrease activity) so fewer enzyme-substrate complexes (ESCs) or less/no substrate can bind/combine/attach/fit into OR (in context of decrease in pH and increase in activity) more ESCs or more substrate binds/attaches;	
	[1]	(in correct context of effect of pH on enzyme i.e. when pH higher/nearer 7/less acidic/more alkaline) <u>denatured/denaturation;</u>	
			[Total: 20]

Question	Expected Answers	Additional guidance	
2 (a) (i)	Draw a large plan diagram showing the features of the wall of the organ. Label the position of the lumen. [4]		
PDO layout 1	[1] Reject if drawn over print of question		
	<table border="1"> <tr> <td>Reject <ul style="list-style-type: none"> thick lines feathery lines one 'tail' or overlap or gap clear, sharp, unbroken lines </td> <td>AND no shading</td> <td>AND uses most of space provided;</td> </tr> </table>		Reject <ul style="list-style-type: none"> thick lines feathery lines one 'tail' or overlap or gap clear, sharp, unbroken lines
Reject <ul style="list-style-type: none"> thick lines feathery lines one 'tail' or overlap or gap clear, sharp, unbroken lines 	AND no shading	AND uses most of space provided;	
MMO collection 2	[1] Reject if drawn two walls		
	<table border="1"> <tr> <td>no cells drawn</td> <td>AND three layers drawn include any circles as only one layer;</td> </tr> </table>		no cells drawn
no cells drawn	AND three layers drawn include any circles as only one layer;		
MMO decisions 1	[1] Reject if only two layers drawn innermost layer is wider than outermost layer at same point;		
MMO decisions 1	[1] Reject <ul style="list-style-type: none"> if any label is biologically incorrect e.g. regions belonging to other organs or plants. label within drawn area – e.g. between two walls correct label with label line to or in lumen on inside;		

(ii) Annotate (make notes with label lines) your diagram to show one difference between the outside layers and the inside layers. [1]					
MMO decision max 1		Reject <ul style="list-style-type: none"> if written over lines of the diagram drawing. if written underneath, unless have labelled on diagram Allow 'er' for one label		max 1	
	[1]	thickness	outermost		innermost
	[1]	Reject cell wall	thin)ner)		think(er);
	[1]	texture	smooth		rough;
	[1]	cells/nuclei	Not clear/densely packed/ visible		Clear/less densely packed/(air) spaces/lots
[1]	Colours/staining of	Pink/red/grey/lighter/more	Purple/darker/less;		
(b) (i) Actual diameter of the nucleus in the cell labelled Y is 7.8 μm. Use this information to calculate the actual diameter of the largest nucleolus in cell Y. [4]					
MMO collection 2	[1]	correct measurement of <u>one</u> nucleus, 11 to 15 mm;		Reject if no units	
	[1]	correct measurement of <u>one</u> nucleolus, 2 to 4.5 mm;		Reject if no units	
PDO display 2	[1]	(mean) adds three measurements	AND shows division by 3;		
	[1]	answer to no more than 2 significant figures, (1 decimal place) between 1.1 and 6.4;		Reject standard form	

(ii) Suggest how you would make the measurement of each nucleolus more accurate.				[1]
ACE improvement 1	[1]	different dimensions/diameters or use vernier callipers or (eyepiece) graticule or increase magnification or high power (of microscope) or enlarging or increase resolution;		
(iii) Make a large drawing of the cell labelled X with three complete cells touching cell X.				[5]
PDO layout 1	[1]	Reject if drawn over print of question Reject <ul style="list-style-type: none"> • thick lines • feathery lines • 2 'tails' or overlaps or gaps 		
		AND no shading	AND uses most of space provided;	
		clear, sharp, unbroken lines		
MMO collection 2	[1]	only cell X and three correct complete touching cells;		
	[1]	nucleus with at least two distinct nucleoli (other than cell X);		
MMO decisions 2	[1]	chromosomes drawn as two areas (no details of chromosomes shown);		
	[1]	blue region/spindle around chromosomes drawn in cell X ;		

(iv) Prepare the space below so that it suitable for you to compare the cells labelled X and Y. [5]						
PDO recording 2	[1]	organise as a table or Venn diagram or ruled connected boxes	headed (cell) <u>X</u> and (cell) <u>Y</u>	differences opposite each other;	<u>X</u> <u>Y</u> 	
	[1]	heading for similarities/similarity/compare (with contrast)/same;				
MMO decision 1	[1]	has at least one correct similarity, cytoplasm or cell/plasma membrane or shape;				
ACE interpretation max 2	Reject tick and cross without a key				<p>if no organisation then mark points only if in same sentence or following sentences.</p> <p>Allow two ticks for both present i.e. for cytoplasm and shape.</p> <p>Allow differences even if not opposite each other.</p> <p>Allow difference on one side if e.g. use more or –er.</p> <p>max 2</p>	
	[1]		feature	(cell) X		(cell) Y
	[1]	1	nucleus/nuclear membrane	absent/none/not clear		present/clear;
	[1]	2	nucleoli	absent/none/		present/clear;
	[1]	3	cytoplasm	less/not granular		more/granular;
	[1]	4	spindle fibres	present/visible		absent/none/not visible;
	[1]	5	chromosomes/chromatid(s)	present/visible		not visible;
	[1]	6	cytoskeleton	absent/not clear		present/clear/visible;
[1]	7	cell size	small(er)	larg(er);		
		Similarities				
[Total: 20]						