

## **MARK SCHEME for the May/June 2013 series**

### **9701 CHEMISTRY**

**9701/31**

Paper 31 (Advanced Practical Skills 1),  
maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9701	31

Question	Sections	Indicative material	Mark
1 (a)	PDO layout	<b>I</b> Constructs a table for results with space for 10 volumes.	1
	PDO recording	<b>II</b> Appropriate headings and units for data given. Volumes in $\text{cm}^3$ or $/\text{cm}^3$ or $(\text{cm}^3)$ , temperature in $^{\circ}\text{C}$ or $/^{\circ}\text{C}$ or $(^{\circ}\text{C})$ in table. All volumes to same dp.	1
	PDO recording	<b>III</b> All temperatures recorded to the nearest $0.5^{\circ}\text{C}$ both in the table and for $T_1$ . At least one ending in .0 and one in .5.	1
	MMO quality	<b>IV + V</b> Compare temp rise for addition of $25\text{ cm}^3$ of <b>FA 2</b> with the Supervisor value. Award 2 marks for $\Delta T$ within $\pm 1^{\circ}\text{C}$ . Award 1 mark for $\Delta T$ within $\pm 2^{\circ}\text{C}$ .	2 [5]
(b)	ACE interpretation	Correctly calculates $\Delta T$ , $V_T$ and $\Delta T \times V_T$ (assume correct data from (a)) (min 8 results)	1 [1]
(c) (i)	PDO layout	<b>I</b> $\Delta T \times V_T$ on <i>y-axis</i> and volume of <b>FA 2</b> on <i>x-axis</i> . Axes clearly labelled (ignore units).	1
		<b>II</b> Uniform scales chosen to use more than half of each axis. Only include 0 if point plotted. Points plotted use 5 large squares vertically and 4 horizontally.	1
		<b>III</b> All points plotted. Examiner to check points at $V = 5, 10, 15, 20$ and $25$ . The points should be within $\frac{1}{2}$ small square and in correct small square. Min 8	1
(c) (ii)		<b>IV</b> Draws both straight lines of best fit.	1
(c) (iii)	ACE interpretation	Reads correctly the value of <b>FA 2</b> from the intercept of the two lines. Answer within $0.5\text{ cm}^3$ . Ignore sf.	1 [5]
(d) (i)		$0.0500\text{ mol}$ (Allow $0.050$ )	1
(d) (ii)		$0.0250\text{ mol}$ (allow $0.025$ ) Allow ecf from (i)/2	1
(d) (iii)		$1000 \times (\text{d})(\text{ii}) / (\text{c})(\text{iii})$ (2–4 sf) Allow ecf from (ii). Penalise sf once only.	1 [3]

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9701	31

(e)	ACE improvements	<p>Accuracy of temperature measurement – use a 0 to 50 °C thermometer or a thermometer with smaller scale divisions (not just more accurate/ electronic thermometer/parallax).</p> <p>Uncertainty about where the lines cross – sample more values of <b>FA 2</b> in the region of the intersection.</p> <p>Repeat/ extra readings on LHS of intersection/ near maximum.</p> <p>Initial temperatures of acid and alkali not same – measure both.</p> <p>Other answers acceptable if <b>specific</b>.</p>	1 [1]
			<b>[Total: 15]</b>
2 (a)	MMO collection	<b>I</b> Initial and final volumes recorded for rough AND initial, final and volume added recorded for accurate titre.	1
	PDO recording	<b>II</b> All accurate readings recorded to 0.05 cm <sup>3</sup> . <i>Do not award if 50(.00) is used as an initial burette reading; more than one final burette reading is 50.(00); any burette reading is greater than 50.(0).</i>	1
	MMO decision	<b>III</b> Two uncorrected accurate titres within 0.1 cm <sup>3</sup> . <i>Do not award if, having performed 2 titres within 0.1 cm<sup>3</sup>, a further titration is performed that is &gt;0.1 cm<sup>3</sup> from the closer of the original 2 titres unless a further titration has been carried out which is within 0.1 cm<sup>3</sup> of any other.</i>	1
	MMO quality	<b>IV + V</b> Award 2 marks if difference from Supervisor within 0.20 cm <sup>3</sup> . Award 1 mark if difference from Supervisor within 0.50 cm <sup>3</sup> .  Examiner compares candidate mean titre with Supervisor mean titre. If best titres are ≥ 0.5 cm <sup>3</sup> , cancel one of the Q marks.	2 [5]

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9701	31

(b)	ACE interpretation	<p>Calculates the mean to appropriate decimal places. <i>The mean should normally be quoted to 2 dp rounded to the nearest 0.01. Example: 26.667 must be rounded to 26.67.</i></p> <p><i>Two special cases where the mean may not be to 2 dp: allow mean to 3 dp only for 0.025 or 0.075 e.g. 26.325; allow mean to 1 dp if <b>all</b> accurate burette readings were given to 1 dp and the mean is exactly correct. eg 26.0 and <math>26.2 = 26.1</math> is correct but 26.0 and <math>26.1 = 26.1</math> is incorrect.</i></p> <p><i>Note: the candidate's mean will sometimes be marked as correct even if it is different from the mean calculated by the Examiner for the purpose of assessing accuracy.</i></p>	1 [1]
(c)	ACE interpretation	<p>All answers correct.</p> <p><b>(i)</b> <math>0.15 \times (\mathbf{b}) / 1000</math></p> <p><b>(ii)</b> <math>(\mathbf{i})/2</math></p> <p><b>(iii)</b> <math>(\mathbf{ii}) \times 400</math></p>	1
	PDO display	Working shown in <b>(i)</b> and <b>(iii)</b>	1
	PDO display	All answers given to 3 or 4 sig figs (minimum 2).	1 [3]
(d)	ACE interpretation	Correctly works out % difference to min 2 sig figs.	1 [1]
<b>[Total: 10]</b>			

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9701	31

FA 5 = HCl    FA 6 = Pb(NO <sub>3</sub> ) <sub>2</sub> FA 7 = K <sub>2</sub> CrO <sub>4</sub> FA 8 = Na <sub>2</sub> CO <sub>3</sub> FA 9 = KBr												
3 (a)	MMO collection	I	FA 5 and FA 6: a white ppt insoluble in excess	1								
		II	FA 5 and FA 7: solution turns from yellow or colourless to orange	1								
		III	FA 5 and FA 8: bubbles or effervescence	1								
	MMO decisions	IV	Uses limewater to test for gas and result.	1								
	ACE conclusion	V	Identifies gas as CO <sub>2</sub> .	1								
	MMO collection	VI	FA 6 and FA 7: a yellow ppt (insol in excess)	1								
		VII	FA 6 and FA 8: a white ppt (insol in excess)	1								
		VIII	FA 7 and FA 8: no reaction/ colourless to yellow (solution)	1 [8]								
(b)	ACE conclusion	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>Pb<sup>2+</sup></td> <td>Cl<sup>-</sup></td> <td>CO<sub>3</sub><sup>2-</sup></td> <td>CrO<sub>4</sub><sup>2-</sup></td> </tr> <tr> <td>FA 6</td> <td>FA 5</td> <td>FA 8</td> <td>FA 7</td> </tr> </table> <p>4 correct scores 2 marks 3 correct scores 1 mark</p>		Pb <sup>2+</sup>	Cl <sup>-</sup>	CO <sub>3</sub> <sup>2-</sup>	CrO <sub>4</sub> <sup>2-</sup>	FA 6	FA 5	FA 8	FA 7	2 [2]
Pb <sup>2+</sup>	Cl <sup>-</sup>	CO <sub>3</sub> <sup>2-</sup>	CrO <sub>4</sub> <sup>2-</sup>									
FA 6	FA 5	FA 8	FA 7									
(c)	ACE conclusion	H <sup>+</sup> because of colour change with chromate or CO <sub>2</sub> / gas released with carbonate.		1 [1]								
(d) (i)	ACE conclusion	No as PbBr <sub>2</sub> / lead bromide is also a white ppt / gives the same observation (if correct in table).		1								
(d) (ii)	MMO decision	Add AgNO <sub>3</sub> followed by NH <sub>3</sub> .		1								
	MMO collection	Cream ppt partially soluble or insoluble in ammonia/ soluble in conc. NH <sub>3</sub> .		1								
	ACE conclusion	<b>Bromide / Br</b> can be allowed from 'off white / buff'.		1 [4]								
<b>[Total: 15]</b>												