





Leave  
blank

**Question 1 continued**

Lined writing area for Question 1.

**(Total 5 marks)**

**Q1**

3

**Turn over**



N 3 5 3 9 0 A 0 3 2 8



Question 2 continued

Leave blank

Handwriting lines for the answer to Question 2.


Q2

(Total 7 marks)



N 3 5 3 9 0 A 0 5 2 8

3.



**Figure 1**

A small box is pushed along a floor. The floor is modelled as a rough horizontal plane and the box is modelled as a particle. The coefficient of friction between the box and the floor is  $\frac{1}{2}$ . The box is pushed by a force of magnitude 100 N which acts at an angle of  $30^\circ$  with the floor, as shown in Figure 1.

Given that the box moves with constant speed, find the mass of the box. (7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---















Question 4 continued

Leave  
blank

Lined area for writing the answer to Question 4.

(Total 7 marks)

Q4



N 3 5 3 9 0 A 0 1 1 2 8









**Question 5 continued**

Leave  
blank

Lined area for writing the answer to Question 5.





6. A ball is projected vertically upwards with a speed of  $14.7 \text{ m s}^{-1}$  from a point which is 49 m above horizontal ground. Modelling the ball as a particle moving freely under gravity, find

(a) the greatest height, above the ground, reached by the ball, **(4)**

(b) the speed with which the ball first strikes the ground, **(3)**

(c) the total time from when the ball is projected to when it first strikes the ground. **(3)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



















8.




Figure 3

Two particles  $A$  and  $B$  have mass  $0.4\text{ kg}$  and  $0.3\text{ kg}$  respectively. The particles are attached to the ends of a light inextensible string. The string passes over a small smooth pulley which is fixed above a horizontal floor. Both particles are held, with the string taut, at a height of  $1\text{ m}$  above the floor, as shown in Figure 3. The particles are released from rest and in the subsequent motion  $B$  does not reach the pulley.

(a) Find the tension in the string immediately after the particles are released. (6)

(b) Find the acceleration of  $A$  immediately after the particles are released. (2)

When the particles have been moving for  $0.5\text{ s}$ , the string breaks.

(c) Find the further time that elapses until  $B$  hits the floor. (9)

---

---

---

---

---

---

---

---

---

---









**Question 8 continued**

Leave  
blank

Lined writing area consisting of 30 horizontal lines.





